

THE CATENARY DEGREE OF KRULL MONOIDS II

ALFRED GEROLDINGER AND QINGHAI ZHONG

ABSTRACT. Let H be a Krull monoid with finite class group G such that every class contains a prime divisor (for example, a ring of integers in an algebraic number field or a holomorphy ring in an algebraic function field). The catenary degree $c(H)$ of H is the smallest integer N with the following property: for each $a \in H$ and each two factorizations z, z' of a , there exist factorizations $z = z_0, \dots, z_k = z'$ of a such that, for each $i \in [1, k]$, z_i arises from z_{i-1} by replacing at most N atoms from z_{i-1} by at most N new atoms. To exclude trivial cases, suppose that $|G| \geq 3$. Then the catenary degree depends only on the class group G and we have $c(H) \in [3, D(G)]$, where $D(G)$ denotes the Davenport constant of G . It is well-known when $c(H) \in \{3, 4, D(G)\}$ holds true. Based on a characterization of the catenary degree determined in the first paper [17], we determine the class groups satisfying $c(H) = D(G) - 1$. Apart from the mentioned extremal cases the precise value of $c(H)$ is known for no further class groups.

1. INTRODUCTION AND MAIN RESULTS

As the title indicates, we continue the investigation of the arithmetic of Krull monoids. All integrally closed noetherian domains are Krull, and holomorphy rings in global fields are Krull monoids with finite class group and infinitely many prime ideals in each class. A Krull monoid is factorial if and only if its class group is trivial, and if this is not the case, then its arithmetic is described by invariants, such as sets of lengths and catenary degrees. We recall some basic definitions.

Let H be a Krull monoid with class group G . Then each non-unit $a \in H$ can be written as a product of atoms, and if $a = u_1 \dots u_k$ with atoms u_1, \dots, u_k of H , then k is called the length of the factorization. The set of lengths $\mathsf{L}(a)$ of all possible factorization lengths is finite, and if $|\mathsf{L}(a)| > 1$, then $|\mathsf{L}(a^n)| > n$ for each $n \in \mathbb{N}$. We denote by $\mathcal{L}(H) = \{\mathsf{L}(a) \mid a \in H\}$ the system of sets of lengths of H . This is an infinite family of finite subsets of positive non-negative integers which is described by a variety of arithmetical parameters. The present paper will focus on the three closely related invariants, namely the set of distances, the $\mathfrak{T}(H)$ invariant, and the catenary degree. If $L = \{m_1, \dots, m_l\} \subset \mathbb{Z}$ is a finite set of integers with $l \in \mathbb{N}$ and $m_1 < \dots < m_l$, then $\Delta(L) = \{m_i - m_{i-1} \mid i \in [2, l]\} \subset \mathbb{N}$ is the set of distances of L . The set of distances $\Delta(H)$ of H is the union of all sets $\Delta(L)$ with $L \in \mathcal{L}(H)$. If

$$\mathfrak{T}(H) = \sup\{\min(L \setminus \{2\}) \mid 2 \in L \in \mathcal{L}(H)\},$$

then $\mathfrak{T}(H) \leq 2 + \sup \Delta(H)$. The catenary degree $c(H)$ of H is defined as the smallest integer N with the following property: for each $a \in H$ and each two factorizations z and z' of a , there exist factorizations $z = z_0, \dots, z_k = z'$ of a such that, for each $i \in [1, k]$, z_i arises from z_{i-1} by replacing at most N atoms from z_{i-1} by at most N new atoms. A simple argument shows that H is factorial if and only if $c(H) = 0$, and if this is not the case, then $2 + \sup \Delta(H) \leq c(H)$.

The study of these arithmetical invariants (in settings ranging from numerical monoids to Mori rings with zero-divisors) has attracted a lot of attention in the recent literature (for a sample see [9, 8, 23, 13, 24, 7, 10]). Our main focus here will be on Krull monoids with finite class group G such that each class contains a

2010 *Mathematics Subject Classification.* 11R27, 11P70, 11B50 13A05, 20M13.

Key words and phrases. non-unique factorizations, sets of lengths, sets of distances, catenary degree, Krull monoids, zero-sum sequences.

This work was supported by the Austrian Science Fund FWF, Project Number M1641-N26.

prime divisor. Let H be such a Krull monoid. Then $|L| = 1$ for all $L \in \mathcal{L}(H)$ if and only if $|G| \leq 2$. Suppose that $|G| \geq 3$. Then the Davenport constant $D(G)$ is finite, $\mathfrak{T}(H) \geq 3$, and there is a canonical chain of inequalities

$$(*) \quad \mathfrak{T}(H) \leq 2 + \max \Delta(H) \leq c(H) \leq D(G).$$

In general, each inequality can be strict (see [17, page 146]). However, for the Krull monoids under consideration the main result in [17] states that $\mathfrak{T}(H) = c(H)$ holds under a certain mild assumption on the Davenport constant. Our starting point is the following Theorem A (the first statement follows from [18, Theorem 6.4.7], and the characterization of $c(H) \in [3, 4]$ is given in [17, Corollary 5.6]).

Theorem A. *Let H be a Krull monoid with finite class group G where $|G| \geq 3$ and each class contains a prime divisor. Then $c(H) \in [3, D(G)]$, and we have*

1. $c(H) = D(G)$ if and only if G is either cyclic or an elementary 2-group.
2. $c(H) = 3$ if and only if G is isomorphic to one of the following groups: $C_3, C_2 \oplus C_2$, or $C_3 \oplus C_3$.
3. $c(H) = 4$ if and only if G is isomorphic to one of the following groups: $C_4, C_2 \oplus C_4, C_2 \oplus C_2 \oplus C_2$, or $C_3 \oplus C_3 \oplus C_3$.

We formulate a main result of the present paper.

Theorem 1.1. *Let H be a Krull monoid with finite class group G where $|G| \geq 3$ and each class contains a prime divisor. Then the following statements are equivalent:*

- (a) $c(H) = D(G) - 1$.
- (b) $\mathfrak{T}(H) = D(G) - 1$.
- (c) G is isomorphic either to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$ or to $C_2 \oplus C_{2n}$ for some $n \geq 2$.

In order to discuss the statements of Theorem 1.1 and their consequences, let H be a Krull monoid as in Theorem 1.1. Then the inequalities in $(*)$ and the fact that $\Delta(H)$ is an interval with $1 \in \Delta(H)$ ([19]) imply that any of the following two conditions,

$$\max \Delta(H) = D(G) - 3 \quad \text{or} \quad \Delta(H) = [1, D(G) - 3]$$

is equivalent to the conditions in Theorem 1.1. The precise value of the Davenport constant is known for p -groups, for groups of rank at most two, and for some others. Thus we do know that $D(C_2^{r-1} \oplus C_4) = r + 3$ and that $D(C_2 \oplus C_{2n}) = 2n + 1$. But the value of $D(G)$ is unknown for general groups of rank three or for groups of the form $G = C_n^r$. Even much less is known for the catenary degree $c(H)$ and for $\mathfrak{T}(H)$. Their precise values are known only for the cases occurring in Theorem A and in Theorem 1.1.

As mentioned at the very beginning, holomorphy rings in global fields are (commutative) Krull monoids with finite class group. In recent years factorization theory has grown towards the non-commutative setting (e.g., [3, 1]) with a focus on maximal orders in central simple algebras (they are non-commutative Krull monoids; see [31, 4]). Combining these results with Theorem 1.1 above, we obtain the following corollary.

Corollary 1.2. *Let \mathcal{O} be a holomorphy ring in a global field K , and R a classical maximal \mathcal{O} -order in a central simple algebra A over K such that every stably free left R -ideal is free. Suppose that the ray class group $G = \mathcal{C}_A(\mathcal{O})$ of \mathcal{O} has at least three elements, let \mathfrak{d} denote a (suitable) distance on R , and $c_{\mathfrak{d}}(R)$ the \mathfrak{d} -catenary degree of R . Then the following statements are equivalent:*

- (a) $c_{\mathfrak{d}}(R) = D(G) - 1$.
- (b) $\mathfrak{T}(R) = D(G) - 1$.
- (c) G is isomorphic either to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$ or to $C_2 \oplus C_{2n}$ for some $n \geq 2$.

Sets of lengths are the most investigated invariants in factorization theory. A standing (but wide open) conjecture states that for the class of Krull monoids under consideration sets of lengths are characteristic. To be more precise, let H and H' be Krull monoids with finite class groups G and G' with $|G| \geq |G'| \geq 4$ and suppose that each class contains a prime divisor. As usual, we write $\mathcal{L}(G) = \mathcal{L}(H)$ and $\mathcal{L}(H') = \mathcal{L}(G')$ (see Proposition 2.1). Then the conjecture states that $\mathcal{L}(G) = \mathcal{L}(G')$ implies that G and G' are isomorphic. For recent work in this direction we refer to [27, 28, 6] or to [18, Section 7.3], [14] for an overview. It turns out the extremal values of $\mathfrak{T}(H)$ discussed in Theorem 1.1 are the main tool to derive an arithmetical characterization of the associated groups. Thus we obtain the following corollary.

Corollary 1.3. *Let G be an abelian group.*

1. *If $\mathcal{L}(G) = \mathcal{L}(C_2^{r-1} \oplus C_4)$ for some $r \geq 2$, then $G \cong C_2^{r-1} \oplus C_4$.*
2. *If $\mathcal{L}(G) = \mathcal{L}(C_2 \oplus C_{2n})$ for some $n \geq 2$, then $G \cong C_2 \oplus C_{2n}$.*

In Section 2 we gather together the required concepts and tools. Section 3 studies sets of lengths in monoids of zero-sum sequences over finite abelian groups, and it is mainly confronted with problems belonging to structural (or inverse) additive number theory. Clearly, the irreducible elements of the monoids are precisely the minimal zero-sum sequences, and we mainly have to deal with minimal zero-sum sequences of extremal length $D(G)$. The structure of minimal zero-sum sequences of length $D(G)$ is known for cyclic groups and elementary 2-groups (in both cases there are trivial answers), and for groups of rank two ([12, 29, 25]). Apart from that, structural results are available only in very special cases ([30, 26]), and this is precisely the lack of information which causes the difficulties in Section 3 (see Prop. 3.5 - 3.8). The proofs of the main results are given in Section 4. They substantially use transfer results (as partly summarized in Proposition 2.1, and also the transfer machinery from [31, 4]), the work from [17] (which relates the catenary degree and the $\mathfrak{T}(\cdot)$ invariant of Krull monoids), and all the work from Section 3.

2. PRELIMINARIES

We denote by \mathbb{N} the set of positive integers and by \mathbb{N}_0 the set of non-negative integers. For $n \in \mathbb{N}$, C_n means a cyclic group of order n . For integers $a, b \in \mathbb{Z}$, $[a, b] = \{x \in \mathbb{Z} \mid a \leq x \leq b\}$ is the discrete interval between a and b . Let $A, B \subset \mathbb{Z}$ be subsets. Then $A + B = \{a + b \mid a \in A, b \in B\}$ denotes their sumset. If $A = \{a_1, \dots, a_l\}$ is finite with $|A| = l \in \mathbb{N}_0$ and $a_1 < \dots < a_l$, then $\Delta(A) = \{a_i - a_{i-1} \mid i \in [2, l]\} \subset \mathbb{N}$ is the set of distances of A . By definition, $\Delta(A) = \emptyset$ if and only if $|A| \leq 1$.

By a monoid, we always mean a commutative semigroup with identity which satisfies the cancellation law (that is, if a, b, c are elements of the monoid with $ab = ac$, then $b = c$ follows). Let H be a monoid. Then H^\times denotes the unit group, $\mathfrak{q}(H)$ the quotient group, $H_{\text{red}} = H/H^\times$ the associated reduced monoid, and $\mathcal{A}(H)$ the set of atoms of H . A monoid F is factorial with $F^\times = \{1\}$ if and only if it is free abelian. If this holds, then the set of primes $P \subset F$ is a basis of F , we write $F = \mathcal{F}(P)$, and every $a \in F$ has a representation of the form

$$a = \prod_{p \in P} p^{v_p(a)} \quad \text{with } v_p(a) \in \mathbb{N}_0 \quad \text{and} \quad v_p(a) = 0 \text{ for almost all } p \in P.$$

If $a \in F$, then $|a| = \sum_{p \in P} v_p(a) \in \mathbb{N}_0$ is the length of a and $\text{supp}(a) = \{p \in P \mid v_p(a) > 0\} \subset P$ is the support of a .

Let G be an additively written abelian group and $G_0 \subset G$ a subset. Then $\langle G_0 \rangle \subset G$ denotes the subgroup generated by G_0 . A family $(e_i)_{i \in I}$ of elements of G is said to be *independent* if $e_i \neq 0$ for all $i \in I$ and, for every family $(m_i)_{i \in I} \in \mathbb{Z}^{(I)}$,

$$\sum_{i \in I} m_i e_i = 0 \quad \text{implies} \quad m_i e_i = 0 \quad \text{for all } i \in I.$$

The family $(e_i)_{i \in I}$ is called a *basis* for G if $G = \bigoplus_{i \in I} \langle e_i \rangle$.

Arithmetical concepts. Our notation and terminology are consistent with [18]. We briefly gather some key notions. The free abelian monoid $Z(H) = \mathcal{F}(\mathcal{A}(H_{\text{red}}))$ is the factorization monoid of H , and the unique homomorphism $\pi: Z(H) \rightarrow H_{\text{red}}$ satisfying $\pi(u) = u$ for all $u \in \mathcal{A}(H_{\text{red}})$ is the factorization homomorphism of H (so π maps a formal product of atoms onto its product in H_{red}). For $a \in H$,

$$\begin{aligned} Z_H(a) = Z(a) = \pi^{-1}(a) \subset Z(H) & \text{ is the set of factorizations of } a, \text{ and} \\ L_H(a) = L(a) = \{|z| \mid z \in Z(a)\} & \text{ is the set of lengths of } a. \end{aligned}$$

Then H is atomic (i.e., each non-unit can be written as a finite product of atoms) if and only if $Z(a) \neq \emptyset$ for each $a \in H$, and H is factorial if and only if $|Z(a)| = 1$ for each $a \in H$. Furthermore, for each $a \in H$, $L(a) = \{0\}$ if and only if $a \in H^\times$, and for all non-units the present definition coincides with the informal one given in the introduction. In particular, $L(a) = \{1\}$ if and only if $a \in \mathcal{A}(H)$. We denote by $\mathcal{L}(H) = \{L(a) \mid a \in H\}$ the system of sets of lengths of H , and by

$$\Delta(H) = \bigcup_{L \in \mathcal{L}(H)} \Delta(L) \quad \text{the set of distances of } H.$$

Distances occurring in sets of lengths L with $2 \in L$ (in other words, in sets of lengths $L(uv)$ with atoms $u, v \in H$) will play a central role. We define

$$\nabla(H) = \sup\{\min(L \setminus \{2\}) \mid L \in \mathcal{L}(H) \text{ with } 2 \in L\}$$

and observe that $\nabla(H) \leq 2 + \sup \Delta(H)$. Before we define catenary degrees we recall the concept of the distance between factorizations. Two factorizations $z, z' \in Z(H)$ can be written in the form

$$z = u_1 \cdots u_l v_1 \cdots v_m \quad \text{and} \quad z' = u_1 \cdots u_l w_1 \cdots w_n$$

with

$$\{v_1, \dots, v_m\} \cap \{w_1, \dots, w_n\} = \emptyset,$$

where $l, m, n \in \mathbb{N}_0$ and $u_1, \dots, u_l, v_1, \dots, v_m, w_1, \dots, w_n \in \mathcal{A}(H_{\text{red}})$. Then $\gcd(z, z') = u_1 \cdots u_l$, and we call $\mathbf{d}(z, z') = \max\{m, n\} = \max\{|z \gcd(z, z')^{-1}|, |z' \gcd(z, z')^{-1}|\} \in \mathbb{N}_0$ the *distance* between z and z' . It is easy to verify that $\mathbf{d}: Z(H) \times Z(H) \rightarrow \mathbb{N}_0$ has all the usual properties of a metric.

Let $a \in H$ and $N \in \mathbb{N}_0 \cup \{\infty\}$. A finite sequence $z_0, \dots, z_k \in Z(a)$ is called an *N -chain of factorizations* if $\mathbf{d}(z_{i-1}, z_i) \leq N$ for all $i \in [1, k]$. We denote by $\mathbf{c}(a)$ the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that any two factorizations $z, z' \in Z(a)$ can be concatenated by an N -chain. Note that $\mathbf{c}(a) \leq \sup L(a)$, that $\mathbf{c}(a) = 0$ if and only if $|Z(a)| = 1$, and if $|Z(a)| > 1$, then $2 + \sup \Delta(L(a)) \leq \mathbf{c}(a)$. Globalizing this concept we define

$$\mathbf{c}(H) = \sup\{\mathbf{c}(b) \mid b \in H\} \in \mathbb{N}_0 \cup \{\infty\}$$

as the *catenary degree* of H . Then $\mathbf{c}(H) = 0$ if and only if H is factorial, and if H is not factorial, then $2 + \sup \Delta(H) \leq \mathbf{c}(H)$.

Krull monoids. A monoid homomorphism $\varphi: H \rightarrow F$ is said to be a divisor homomorphism if $\varphi(a) \mid \varphi(b)$ in F implies that $a \mid b$ in H for all $a, b \in H$. A monoid H is said to be a Krull monoid if one of the following equivalent properties is satisfied (see [18, Theorem 2.4.8] or [21]):

- (a) H is completely integrally closed and satisfies the ascending chain condition on divisorial ideals.
- (b) H has a divisor homomorphism into a free abelian monoid.
- (c) H has a divisor theory: this is a divisor homomorphism $\varphi: H \rightarrow F = \mathcal{F}(P)$ into a free abelian monoid such that for each $p \in P$ there is a finite set $E \subset H$ with $p = \gcd(\varphi(E))$.

Let H be a Krull monoid. Then a divisor theory is unique up to unique isomorphism and the group $\mathcal{C}(\varphi) = \mathfrak{q}(F)/\mathfrak{q}(\varphi(H))$ depends only on H , and hence it is called the class group of H . We say that a class $g = [a] = a\mathfrak{q}(\varphi(H)) \subset \mathfrak{q}(F) \in \mathcal{C}(\varphi)$, with $a \in \mathfrak{q}(F)$, contains a prime divisor if $g \cap P \neq \emptyset$. A domain R is Krull if and only if its multiplicative monoid $R^\bullet = R \setminus \{0\}$ of non-zero elements is Krull. Thus Property (a) shows that every integrally closed noetherian domain is Krull. If R is Krull with finite class group such that each class contains a prime divisor, then the same is true for regular congruence submonoids of R ([18, Section 2.11]). For monoids of modules which are Krull we refer the reader to [5, 2, 11].

Next we discuss a Krull monoid having a combinatorial flavor. It plays a universal role in all arithmetical studies of general Krull monoids. Let G be an additive abelian group and $G_0 \subset G$ a subset. According to the tradition in combinatorial number theory, elements $S \in \mathcal{F}(G_0)$ will be called sequences over G_0 (for a recent presentation of their theory we refer to [20], and for an overview of their interplay with factorization theory we refer to [14]). Let $S = g_1 \cdot \dots \cdot g_l = \prod_{g \in G_0} g^{v_g(S)} \in \mathcal{F}(G_0)$ be a sequence over G_0 . Then $\sigma(S) = g_1 + \dots + g_l \in G$ is the sum of S and

$$\Sigma(S) = \left\{ \sum_{i \in I} g_i \mid \emptyset \neq I \subset [1, l] \right\} = \left\{ \sigma(T) \mid T \text{ is a subsequence of } S \right\} \subset G$$

denotes the set of subsums of S . We say that S is zero-sum free if $0 \notin \Sigma(S)$ and that S is a zero-sum sequence if $\sigma(S) = 0$. Obviously, S is zero-sum free or a (minimal) zero-sum sequence if and only if $-S = (-g_1) \cdot \dots \cdot (-g_l)$ has this property. The set

$$\mathcal{B}(G_0) = \{S \in \mathcal{F}(G_0) \mid \sigma(S) = 0\} \subset \mathcal{F}(G_0)$$

of all zero-sum sequences over G_0 is a submonoid of $\mathcal{F}(G_0)$, and since the embedding $\mathcal{B}(G_0) \hookrightarrow \mathcal{F}(G_0)$ obviously is a divisor homomorphism, Property (b) shows that $\mathcal{B}(G_0)$ is a Krull monoid. For each arithmetical invariant $*(H)$ defined for a monoid H , we write $*(G_0)$ instead of $*(\mathcal{B}(G_0))$. This is the usual convention and will hardly lead to misunderstandings. In particular, we set $\mathcal{L}(G_0) = \mathcal{L}(\mathcal{B}(G_0))$, $\mathfrak{T}(G_0) = \mathfrak{T}(\mathcal{B}(G_0))$, and $\Delta(G_0) = \Delta(\mathcal{B}(G_0))$. The set $\mathcal{A}(G_0) = \mathcal{A}(\mathcal{B}(G_0))$ of atoms of $\mathcal{B}(G_0)$ is the set of minimal zero-sum sequences over G_0 and

$$D(G_0) = \sup\{|A| \mid A \in \mathcal{A}(G_0)\} \in \mathbb{N} \cup \{\infty\}$$

is the Davenport constant of G_0 . If G_0 is finite, then $\mathcal{A}(G_0)$ is finite and hence $D(G_0) < \infty$.

Suppose that G is finite abelian, say $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$ with $1 < n_1 \mid \dots \mid n_r$. Then $1 + \sum_{i=1}^r (n_i - 1) \leq D(G)$. We will use without further mention that equality holds for p -groups and for groups of rank $r \leq 2$ ([18, Chapter 5]). Furthermore, we will frequently use that, if $\exp(G) + 1 = n_r + 1 = D(G)$, then $r = 2$ and $G \cong C_2 \oplus C_{n_2}$. If $S \in \mathcal{F}(G)$ is zero-sum free of length $|S| = D(G) - 1$, then $\Sigma(S) = G \setminus \{0\}$, and if $S \in \mathcal{A}(G)$ with $|S| = D(G)$, then $\Sigma(S) = G$ (see [18, Proposition 5.1.4]).

Suppose that $|G| \geq 3$. Then $\mathcal{B}(G)$ is a Krull monoid whose class group is isomorphic to G and each class contains precisely one prime divisor ([18, Proposition 2.5.6]). Furthermore, its arithmetic reflects the arithmetic of more general Krull monoids as it is summarized in the next proposition (for a proof see [18, Section 3.4]).

Proposition 2.1. *Let H be a Krull monoid, $\varphi: H \rightarrow F = \mathcal{F}(P)$ a divisor theory, $G = \mathcal{C}(\varphi)$ the class group, and suppose that each class contains a prime divisor. Let $\tilde{\beta}: \mathcal{F}(P) \rightarrow \mathcal{F}(G)$ denote the unique homomorphism satisfying $\tilde{\beta}(p) = [p]$ for each $p \in P$, and let $\beta = \tilde{\beta} \circ \varphi: H \rightarrow \mathcal{B}(G)$.*

1. *For each $a \in H$, we have $\mathsf{L}_H(a) = \mathsf{L}_{\mathcal{B}(G)}(\beta(a))$. In particular, $\mathcal{L}(H) = \mathcal{L}(G)$, $\Delta(H) = \Delta(G)$, and $\mathfrak{T}(H) = \mathfrak{T}(G)$.*
2. *We have $|G| \leq 2$ if and only if $D(G) \leq 2$ if and only if $|L| = 1$ for all $L \in \mathcal{L}(G)$.*
3. *If $|G| \geq 3$, then $\mathsf{c}(H) = \mathsf{c}(G)$.*

3. ON SETS OF LENGTHS $L \in \mathcal{L}(G)$ HAVING EXTREMAL PROPERTIES

In this section we mainly study sets of lengths of zero-sum sequences over finite abelian groups. We start by recalling two results from [17].

Lemma 3.1. *Let G be a finite abelian group with $|G| \geq 3$, say $G = C_{n_1} \oplus \dots \oplus C_{n_r}$ with $1 < n_1 \mid \dots \mid n_r$.*

1. $\mathsf{c}(G) \leq \max \left\{ \left\lfloor \frac{1}{2} D(G) + 1 \right\rfloor, \mathfrak{T}(G) \right\}$.
2. $\mathfrak{T}(G) \geq \max \left\{ n_r, 1 + \sum_{i=1}^r \left\lfloor \frac{n_i}{2} \right\rfloor \right\}$.

Proof. See Proposition 4.1 and Theorem 4.2 in [17]. \square

Lemma 3.2. *Let G be an abelian group with $|G| \geq 3$, and let $U, V \in \mathcal{A}(G)$ with $\max \mathbf{L}(UV) \geq 3$.*

1. *Let $K \subset G$ be a finite cyclic subgroup. If $\sum_{h \in K} \mathbf{v}_h(UV) > |K|$ and there exists a non-zero $g \in K$ such that $\mathbf{v}_g(U) > 0$ and $\mathbf{v}_{-g}(V) > 0$, then $\mathbf{L}(UV) \cap [3, |K|] \neq \emptyset$.*
2. *If $\mathbf{L}(UV) \cap [3, \text{ord}(g)] = \emptyset$, then for any $g \in G$, we have $\mathbf{v}_g(U) + \mathbf{v}_{-g}(V) \leq \text{ord}(g)$.*

Proof. 1. See Lemma 5.2 in [17].

2. This follows from 1. with $K = \langle g \rangle$. \square

Let A be a zero-sum sequence over a finite abelian group G and suppose that $0 \nmid A$. If

$$A = U_1 \cdot \dots \cdot U_k = V_1 \cdot \dots \cdot V_l,$$

where $k, l \in \mathbb{N}$ and $U_1, \dots, U_k, V_1, \dots, V_l \in \mathcal{A}(G)$, then obviously

$$2l \leq \sum_{i=1}^l |V_i| = |A| = \sum_{i=1}^k |U_i| \leq k\mathbf{D}(G).$$

These inequalities will be used implicitly in many of the forthcoming arguments.

Lemma 3.3. *Let $G = C_4 \oplus C_4$. Then $\mathfrak{T}(G) = \mathbf{c}(G) = 5$. Moreover, if $U, V \in \mathcal{A}(G)$ with $\mathbf{L}(UV) \cap [2, 5] = \{2, 5\}$, then $\langle \text{supp}(UV) \rangle = G$.*

Proof. First, we prove the moreover statement. Let $U, V \in \mathcal{A}(G)$ with $\mathbf{L}(UV) \cap [2, 5] = \{2, 5\}$. Then $\mathfrak{T}(\langle \text{supp}(UV) \rangle) \geq 5$. Since for every proper subgroup K of G we have $\mathfrak{T}(K) \leq \mathfrak{T}(C_2 \oplus C_4) < \mathbf{D}(G) = 5$ by Theorem A, it follows that $\langle \text{supp}(UV) \rangle = G$.

Recall that $\mathbf{D}(G) = 7$, and note that it suffices to prove $\mathfrak{T}(G) \leq 5$, since then combining with [17, Proposition 4.1.2 and Corollary 4.3] yields $5 \leq \mathfrak{T}(G) = \mathbf{c}(G) \leq 5$.

Let $U, V \in \mathcal{A}(G)$ with $\max \mathbf{L}(UV) > 5$ be given. Since $\max \mathbf{L}(UV) \leq \min\{|U|, |V|\} \leq \mathbf{D}(G) = 7$, it follows that $\min\{|U|, |V|\} \in \{6, 7\}$. We have to show that there exists a factorization $UV = W_1 \cdot \dots \cdot W_k$ with $W_1, \dots, W_k \in \mathcal{A}(G)$ and $k \in [3, 5]$. We start with two special cases.

First, suppose that $V = -U$ and $|U| = 7$. Then [18, Theorem 6.6.7] implies that $4 \in \mathbf{L}(UV)$, and thus the assertion follows.

Second, suppose that there exist $W_1, W_2 \in \mathcal{A}(G)$ such that $W_1 W_2 \mid UV$, $5 \geq |W_1| \geq |W_2|$, and $|W_1 W_2| \geq 7$. Then there exist $k \in \mathbb{N}_{\geq 3}$, $W_3, \dots, W_k \in \mathcal{A}(G)$ such that $UV = W_1 \cdot \dots \cdot W_k$, and

$$2(k-2) \leq |W_3 \cdot \dots \cdot W_k| = |UV| - |W_1 W_2| \leq 14 - 7 = 7$$

implies $k \leq 5$, and the assertion follows.

Assume to the contrary that UV has no factorization of length $k \in [3, 5]$. Then none of the two special cases holds true. By [16, Lemma 3.6], UV has a zero-sum subsequence $W_1 \in \mathcal{A}(G)$ of length $|W_1| \in [2, 4]$, and suppose that $|W_1|$ is maximal. Then there is a factorization

$$UV = W_1 \cdot \dots \cdot W_k \quad \text{with } k \geq 3 \quad \text{and } W_1, \dots, W_k \in \mathcal{A}(G).$$

By assumption we have $k \geq 6$. Since $k = 7$ would imply that $V = -U$ and $|U| = 7$, it follows that $k = 6$. We distinguish three cases.

CASE 1: $|W_1| = 2$.

Since $|W_1|$ is maximal, we get $|W_1| = \dots = |W_k| = 2$, and thus $|UV| \in \{12, 14\}$ and $V = -U$. Since we are not in the first special case, it follows that $|U| = 6$, say $U = g_1 \cdot \dots \cdot g_6$.

If $\mathbf{h}(U) = 3$, say $g_1 = g_2 = g_3$, then $W = (-g_1)g_4g_5g_6 \in \mathcal{A}(G)$ with $W \mid UV$, a contradiction. Thus $\mathbf{h}(U) \leq 2$. Since $\mathbf{D}(C_2^2) = 3$, $\text{supp}(U)$ contains at most two elements of order 2, say $\text{ord}(g_1) = \dots = \text{ord}(g_4) = 4$. Since $\mathbf{D}(C_2 \oplus C_4) = 5$, it follows that $\text{supp}(U)$ generates G . Recall that the maximal size of

a minimal generating set of G equals $r^*(G) = 2$, and that every generating set contains a basis (see [18, Appendix A]). Thus the elements of order 4 contain a basis.

Suppose there are two elements with multiplicity two, say $g_1 = g_3$ and $g_2 = g_4$. Then (g_1, g_2) is a basis of G and $g_5 = ag_1 + bg_2$ with $a, b \in [1, 3]$. Thus there is a zero-sum subsequence $W \in \mathcal{A}(G)$ with $|W| > 2$, $W \mid UV$ and $\text{supp}(W) \in \{g_5, g_1, g_2, -g_1, -g_2\}$, a contradiction.

Suppose there is precisely one element with multiplicity two, say $g_1 = g_3$. Then (g_1, g_2) is a basis of G and there is an element in $\text{supp}(U)$ of the form $ag_1 + bg_2$ with $a \in [1, 3]$ and $b \in \{1, 3\}$. As above we obtain a zero-sum sequence W with $|W| > 2$, a contradiction.

Suppose that $h(U) = 1$. Then (g_1, g_2) is a basis of G . Then there is one element in $\text{supp}(U)$ which is not of the form $\{g_1 + 2g_2, 2g_1 + g_2, 2g_1 + 2g_2\}$, and hence it has the form $ag_1 + bg_2$ with $a, b \in \{1, 3\}$, and we obtain a contradiction as above. \square

CASE 2: $|W_1| = 4$.

Since $k = 6$, $|UV| \leq 14$ and $|W_1| = 4$, we get $|W_2| = \dots = |W_6| = 2$ and $|U| = |V| = 7$. By [18, Example 5.8.8], there exists a basis (e_1, e_2) of G such that

$$U = e_1^3 \prod_{\nu=1}^4 (a_\nu e_1 + e_2)$$

with $a_1, a_2, a_3, a_4 \in [0, 3]$ and $a_1 + a_2 + a_3 + a_4 \equiv 1 \pmod{4}$. We set $X = \gcd(W_1, U)$. Then there are $X, Y, Z \in \mathcal{F}(G)$ such that $U = XZ$, $V = Y(-Z)$ and $W_1 = XY$. After renumbering if necessary there are the following three cases:

$$X = e_1^2, \quad X = e_1(a_1e_1 + e_2) \quad \text{or} \quad X = (a_1e_1 + e_2)(a_2e_1 + e_2).$$

If $X = e_1^2$, then $(-e_1)(a_1e_1 + e_2) \cdot \dots \cdot (a_4e_1 + e_2)$ is a minimal zero-sum subsequence of UV of length 5, a contradiction.

Suppose that $X = (a_1e_1 + e_2)(a_2e_1 + e_2)$. If $a_3 \neq a_4$, then $(a_3e_1 + e_2)(-a_4e_1 - e_2)e_1^3$ has a zero-sum subsequence of length 4 or 5, a contradiction. Suppose that $a_3 = a_4$. Then $(e_1, a_3e_1 + e_2)$ is a basis, and after changing notation if necessary we may suppose that $a_3 = 0$. Then $(a_1e_1 + e_2)(a_2e_1 + e_2) \in \{e_1(e_1 + e_2), (2e_1 + e_2)(-e_1 + e_2)\}$. In the first case $e_1^3(e_1 + e_2)(-e_2)$ and in the second case $(-e_1)^3(-e_1 + e_2)(-e_2)$ is a zero-sum subsequence of UV of length 5, a contradiction.

Suppose that $X = e_1(a_1e_1 + e_2)$. If two of the a_2, a_3, a_4 are distinct, say $a_2 \neq a_3$, then $(a_2e_1 + e_2)(-a_3e_1 - e_2)e_1^2(-e_1)^2$ has a zero-sum subsequence of length greater than 2, a contradiction. Suppose that $a_2 = a_3 = a_4$. Then $(e_1, e_2' = a_2e_1 + e_2)$ is a basis. Then $U = e_1^3(e_1 + e_2')e_2'^3$ and $e_1^3(e_1 + e_2')(-e_2')$ is a minimal zero-sum subsequence of UV of length 5, a contradiction.

CASE 3: $|W_1| = 3$.

After renumbering if necessary we may suppose that $|U| = 7$, $|V| \in \{6, 7\}$, $|W_2| \in \{2, 3\}$ and $|W_3| = \dots = |W_6| = 2$. By [18, Example 5.8.8], there exists a basis (e_1, e_2) of G such that

$$U = e_1^3 \prod_{\nu=1}^4 (a_\nu e_1 + e_2)$$

with $a_1, a_2, a_3, a_4 \in [0, 3]$ and $a_1 + a_2 + a_3 + a_4 \equiv 1 \pmod{4}$. Thus U has a subsequence S of length $|S| = 4$ such that $-S$ is a subsequence of V .

Suppose there are two distinct elements in $\{a_1e_1 + e_2, \dots, a_4e_1 + e_2\}$, say $(a_1e_1 + e_2)$ and $(a_2e_1 + e_2)$ such that $(a_1e_1 + e_2)(a_2e_1 + e_2) \mid S$. Then either $(a_1e_1 + e_2)(-a_2e_1 - e_2)e_1^3$ or $(-a_1e_1 - e_2)(a_2e_1 + e_2)e_1^3$ contains a zero-sum subsequence of length 4 or 5, a contradiction.

Now we suppose that this does not hold and distinguish three cases.

Suppose $S = e_1(a_1e_1 + e_2)(a_2e_1 + e_2)(a_3e_1 + e_2)$. Then $a_1 = a_2 = a_3 = a$, and since $(e_1, ae_1 + e_2)$ is a basis of G , we may suppose that $a = 0$. Then $U = e_1^3e_2^3(e_1 + e_2)$ and $(-e_1)e_2^3(e_1 + e_2)$ is a zero-sum subsequence of UV of length 5, a contradiction.

Suppose $S = e_1^2(a_1e_1 + e_2)(a_2e_1 + e_2)$. Then $a_1 = a_2$, and since $(e_1, ae_1 + e_2)$ is a basis of G , we may suppose that $a = 0$. Then $U = e_1^3e_2^2(ae_1 + e_2)((1-a)e_1 + e_2)$, and either $(-e_2)e_1^3(ae_1 + e_2)$ or $(-e_2)e_1^3((1-a)e_1 + e_2)$ has a zero-sum subsequence of length greater than or equal to 4, a contradiction.

Suppose $S = e_1^3(a_1e_1 + e_2)$. Again we may suppose that $a = 0$ and find a zero-sum subsequence of UV of length greater than or equal to 4, a contradiction. \square

Lemma 3.4. *Let G be a finite abelian group with $|G| \geq 3$ and $U \in \mathcal{A}(G)$ with $|U| = \mathsf{D}(G)$.*

1. *If $g_1, g_2, h \in \text{supp}(U)$, then $g_1 \neq 2g_2$, and if $\text{ord}(h) = 2$, then $g_1 \neq 2g_2 + h$.*
2. *Let $g \in G$ with $\nu_g(U) = k \geq 1$ and $\text{ord}(g) > 2$ and suppose that $|\text{supp}(U)| \geq 2$. Then there exists some $W \in \mathcal{A}(G)$ such that $W | (-g)^k g^{-k}U$ and $|W| \in [3, \mathsf{D}(G) - 1]$.*

Proof. 1. Assume to the contrary that $g_1 = 2g_2$. Since $|U| = \mathsf{D}(G)$, it follows that $\Sigma(g_1^{-1}U) = G \setminus \{0\}$. Hence there exists some $W \in \mathcal{F}(G)$ such that $W | g_1^{-1}U$ and $\sigma(W) = -g_2$. If $g_2 | W$, then $g_1g_2^{-1}W$ is a proper zero-sum subsequence of U , a contradiction. If $g_2 \nmid W$, then g_2W is a proper zero-sum subsequence of U , a contradiction.

Assume to the contrary that $\text{ord}(h) = 2$ and $g_1 = 2g_2 + h$. There exists some $W \in \mathcal{F}(G)$ such that $W | g_1^{-1}U$ and $\sigma(W) = -g_2 + h$. If $g_2 | W$, then $g_1g_2^{-1}W$ is a proper zero-sum subsequence of U , a contradiction. If $g_2 \nmid W$ and $h | W$, then $g_2h^{-1}W$ is a proper zero-sum subsequence of U , a contradiction. If $g_2 \nmid W$ and $h \nmid W$, then g_2hW is a proper zero-sum subsequence of U , a contradiction.

2. Since $|(-g)^k g^{-k}U| = |U| = \mathsf{D}(G)$, there exists some $W \in \mathcal{A}(G)$ such that $W | (-g)^k g^{-k}U$. It is easy to show that $|W| \neq 2$. Assume the contrary that $|W| = \mathsf{D}(G)$. Then $W = (-g)^k g^{-k}U$ and $\text{ord}(g) = 2k > k + 1$. Let $h \in \text{supp}(U) \setminus \{g\}$. Since $\Sigma(h^{-1}U) = G \setminus \{0\}$, there exists some $T \in \mathcal{F}(G)$ such that $T | h^{-1}U$ and $\sigma(T) = (k+1)g$. If $g \nmid T$, then $g^{k-1}T$ is a proper zero-sum subsequence of U , a contradiction. Suppose that $g | T$, say $T = g^t T_0$ with $t \in [1, k]$ and $T_0 | g^{-k}U$. Then $W' = (-g)^{k-t+1}T_0$ is a zero-sum subsequence of $W = (-g)^k g^{-k}U$. Since W is an atom, it follows that $W' = W$. This implies that $T_0 = g^{-k}U$, a contradiction to $T_0 | T | h^{-1}U$. \square

Proposition 3.5. *Let G be a finite abelian group with $\mathsf{D}(G) \geq 5$. Then the following statements are equivalent:*

- (a) *G is isomorphic to $C_2 \oplus C_{2n}$ with $n \geq 2$.*
- (b) *There exist $U, V \in \mathcal{A}(G)$ with $\mathsf{L}(UV) = \{2, \mathsf{D}(G) - 1, \mathsf{D}(G)\}$.*

Proof. (a) \Rightarrow (b) Let (e_1, e_2) be a basis of G with $\text{ord}(e_1) = 2$ and $\text{ord}(e_2) = 2n$ with $n \geq 2$. Then $\mathsf{D}(G) = 2n + 1$, $U = e_1e_2^{2n-1}(e_1 + e_2) \in \mathcal{A}(G)$, and $\mathsf{L}(U(-U)) = \{2, \mathsf{D}(G) - 1, \mathsf{D}(G)\}$.

(b) \Rightarrow (a) Since $\mathsf{D}(G) \in \mathsf{L}(UV)$, it follows that $|U| = |V| = \mathsf{D}(G)$ and $V = -U$. Let $A \in \mathcal{A}(G)$ with $A | U(-U)$. Then $|A| \in \{2, 3, \mathsf{D}(G)\}$, and if $|A| = \mathsf{D}(G)$, then $U(-U)A^{-1}$ is an atom of length $\mathsf{D}(G)$. Since $|U| = \mathsf{D}(G)$, it follows that $\Sigma(U) = G$. By [18, Theorem 6.6.3], G is neither cyclic nor an elementary 2-group. Therefore, $|\text{supp}(U)| \geq 2$ and may write $U = g_1^{k_1} \cdots g_s^{k_s}$ with $g_1, \dots, g_s \in G$ pairwise distinct, $s \geq 2$, $k_1, \dots, k_s \in \mathbb{N}$, and $\text{ord}(g_1) > 2$.

We continue with the following assertion.

- A1.** If $\text{ord } g_i > 2$ for some $i \in [1, s]$, then there exist distinct elements $h_1, h_2 \in \text{supp}(U)$ such that $g_i = h_1 + h_2$.
- A2.** If $\text{ord } g_i > 2$ for some $i \in [1, s]$, then there exist distinct elements $f_1, f_2 \in \text{supp}(U)$ such that $g_i = f_1 + f_2$, $\text{ord}(f_1) > 2$, and $\text{ord}(f_2) = 2$.

Proof of A1. Let $i \in [1, s]$ with $\text{ord}(g_i) > 2$. By Lemma 3.4.2, there exists an atom $W \in \mathcal{A}(G)$ such that $W | g_i^{-k_i}(-g_i)^{k_i}U$ and $|W| \in [3, \mathsf{D}(G) - 1]$. Then $W | U(-U)$ infers $|W| = 3$. Thus $W = (-g_i)^2 h_1$ with $h_1 | U$ or $W = (-g_i)h_1h_2$ with $h_1h_2 | U$. In the first case $h_1 = 2g_i$, a contradiction to Lemma 3.4.1. Thus the second case holds, and again Lemma 3.4.1 implies that h_1 and h_2 are distinct. \square

Proof of A2. Let $i \in [1, s]$ with $\text{ord}(g_i) > 2$, say $i = 1$. By **A1**, we may assume that $g_1 = g_2 + g_3$ and hence $\text{ord}(g_2) > 2$ or $\text{ord}(g_3) > 2$. Assume to the contrary that $\text{ord}(g_2) > 2$ and $\text{ord}(g_3) > 2$. Again by **A1**, there are distinct $h_1, h_2 \in \text{supp}(U)$ such that $g_2 = h_1 + h_2$. Then $\{h_1, h_2\} \cap \{g_1, g_3\} \neq \emptyset$ otherwise $(-g_1)h_1h_2g_3$ would be an atom of length 4 dividing $U(-U)$. Since $g_2 = g_1 + h_i$ with $i \in [1, 2]$ cannot hold, we obtain that $g_2 = g_3 + h$ with $h \in \{h_1, h_2\}$. Repeating the argument we infer that $g_3 = g_2 + h'$ with $h' \in \text{supp}(U)$. It follows that $h + h' = 0$ which implies that $h = h'$, $\text{ord}(h) = 2$, and $g_1 = 2g_3 + h$, a contradiction to Lemma 3.4.1. \square

Since $\text{ord}(g_1) > 2$, by **A2** we may assume that $g_1 = g_2 + g_3$ with $\text{ord}(g_2) > 2$ and $\text{ord}(g_3) = 2$. If $s = 3$, then $G = \langle \text{supp}(U) \rangle = \langle g_1, g_2, g_3 \rangle = \langle g_2, g_3 \rangle \cong C_2 \oplus C_{2n}$ with $n \geq 2$. Assume to the contrary that $s \geq 4$. We distinguish two cases.

CASE 1: There is an $i \in [4, s]$ such that $\text{ord}(g_i) > 2$, say $i = 4$.

By **A2**, we may assume $g_4 = h_1 + h_2$ with $\text{ord}(h_1) > 2$ and $\text{ord}(h_2) = 2$. We assert that $\{g_1, g_2, g_3\} \cap \{g_4, h_1, h_2\} = \emptyset$. Assume to the contrary that this does not hold. Taking into account the order of the elements and that $|\{g_1, \dots, g_4\}| = 4$ we have to consider the following three cases:

- If $h_2 = g_3$, then $g_1 + g_4 = g_2 + h_1$ and hence $(-g_1)(-g_4)g_2h_1$ is an atom of length 4 dividing $U(-U)$, a contradiction.
- If $h_1 = g_1$, then $(-g_4)g_2g_3h_2$ is an atom of length 4 dividing $U(-U)$, a contradiction.
- If $h_1 = g_2$, then $(-g_4)g_1g_3h_2$ is an atom of length 4 dividing $U(-U)$, a contradiction.

Thus we have $\{g_1, g_2, g_3\} \cap \{g_4, h_1, h_2\} = \emptyset$. Obviously, $(-g_1)g_2g_3$, $g_1(-g_2)(-g_3)$, $(-g_4)h_1h_2$, and $g_4(-h_1)(-h_2)$ are atoms of length 3 dividing $U(-U)$, and therefore their product $(-g_1)g_2g_3 \cdot g_1(-g_2)(-g_3) \cdot (-g_4)h_1h_2 \cdot g_4(-h_1)(-h_2)$ divides $U(-U)$, a contradiction to $\mathsf{L}(U(-U)) = \{2, \mathsf{D}(G) - 1, \mathsf{D}(G)\}$.

CASE 2: $\text{ord}(g_i) = 2$ for all $i \in [4, s]$.

If $k_1 \geq 2$ and $k_2 \geq 2$, then $(-g_1)^2g_2^2$ is an atom of length 4 dividing $U(-U)$, a contradiction. Thus $k_1 = 1$ or $k_2 = 1$, say $k_1 = 1$. Since $g_1 + k_2g_2 + g_3 + \dots + g_s = \sigma(U) = 0$ and $(k_2 + 1)g_2 = g_4 + \dots + g_s$, it follows that $\text{ord}(g_2) = 2(k_2 + 1)$. Thus

$$\exp(G) + s - 3 \leq \mathsf{D}(G) = |U| = k_2 + s - 1 \leq 2(k_2 + 1) + s - 4 \leq \exp(G) + s - 4,$$

a contradiction. \square

Proposition 3.6. *Let G be a finite abelian group with $\mathsf{D}(G) \geq 5$. Then there are no $U, V \in \mathcal{A}(G)$ with $\mathsf{L}(UV) = \{2, \mathsf{D}(G) - 1\}$ and $|U| = |V| = \mathsf{D}(G)$.*

Proof. Assume to the contrary that $U, V \in \mathcal{A}(G)$ with these properties do exist. If $A \in \mathcal{A}(G)$ with $A | UV$, then $|A| \in \{2, 3, 4, \mathsf{D}(G)\}$. Since $\Sigma(U) = \Sigma(V) = G$ and G cannot be cyclic, it follows that $|\text{supp}(U)| \geq 2$ and $|\text{supp}(V)| \geq 2$. We distinguish two cases.

CASE 1: For all $A \in \mathcal{A}(G)$ with $A | UV$ we have that $|A| \in \{2, 3, \mathsf{D}(G)\}$.

Then UV has a factorization of the form $UV = W_1 \cdot \dots \cdot W_{\mathsf{D}(G)-1}$ where $W_1, \dots, W_{\mathsf{D}(G)-1} \in \mathcal{A}(G)$, $|W_{\mathsf{D}(G)-2}| = |W_{\mathsf{D}(G)-1}| = 3$, and all the other W_i have length 2. Thus we may set

$$U = g_1 \cdot \dots \cdot g_l h_1 h_4 h_5 \quad \text{and} \quad V = (-g_1) \cdot \dots \cdot (-g_l)(-h_2)(-h_3)(-h_6),$$

where $l \in \mathbb{N}$, $g_1, \dots, g_l, h_1, \dots, h_6 \in G$ not necessarily distinct such that $h_1 = h_2 + h_3$, $h_6 = h_4 + h_5$ and $\{h_1, h_4, h_5\} \cap \{h_2, h_3, h_6\} = \emptyset$.

Since $V(-g_6)^{-1}$ is a zero-sum free sequence of length $\mathsf{D}(G) - 1$, there exists a subsequence T of $V(-g_6)^{-1}$ such that $\sigma(T) = h_4$. Thus $Th_5(-h_6)$ is a zero-sum subsequence of UV . Since $T(-h_6)$ is zero-sum free, we obtain that $Th_5(-h_6)$ is an atom of length $|Th_5(-h_6)| \in \{3, \mathsf{D}(G)\}$. If $|Th_5(-h_6)| = 3$, then $|T| = 1$ which implies that $h_4 | V$. If $|Th_5(-h_6)| = \mathsf{D}(G)$, then $Th_5(-h_6) = V$ which implies that $h_5 | V$. Therefore, we obtain that $h_4 \in \text{supp}(U) \cap \text{supp}(V)$ or $h_5 \in \text{supp}(U) \cap \text{supp}(V)$. Without loss of generality, we assume that $h_4 \in \text{supp}(U) \cap \text{supp}(V)$.

We start with the following assertions.

A1. $\text{ord}(h_4) > 2$, $(-2h_4)^2 \mid U$, and $(-2h_4)^2 \mid V$.

A2. If $g_i \notin \{h_1, \dots, h_6\}$ for some $i \in [1, l]$, then $\text{ord}(g_i) = 2$.

Proof of A1. Since $-g_i \neq h_4$ for all $i \in [1, l]$, we obtain that $h_4 \in \{-h_2, -h_3, -h_6\}$ which implies that $\text{ord}(h_4) > 2$.

By the symmetry of U and V , we only need to prove that $(-2h_4)^2 \mid U$. For any $g \in \text{supp}(U)$ with $g \neq h_4$, consider the sequence $Ug^{-1}h_4$. Since $|Ug^{-1}h_4| = \text{D}(G)$ and $Ug^{-1}h_4 \mid UV$, there exists an atom A with $A \mid Ug^{-1}h_4$ and $|A| = 3$ by Lemma 3.4.2. Note that $h_4^2 \mid A$. Therefore, $-2h_4 \mid Ug^{-1}$. If $\text{ord}(h_4) = 3$, then $h_4 = -2h_4$ and $A = h_4^3$ which implies that $h_4^2 \mid U$. If $\text{ord}(g_4) > 3$. Then $-2h_4 \neq h_4$. Thus we can choose $g = -2h_4$ which implies that $(-2h_4)^2 \mid U$. \square

Proof of A2. Let $i \in [1, l]$ such that $g_i \notin \{h_1, \dots, h_6\}$ and assume to the contrary that $\text{ord}(g_i) > 2$. Let $v = \nu_{g_i}(U) = \nu_{-g_i}(V)$. By Lemma 3.4.2, we obtain that there exists an atom W such that $W \mid g_i^{-v}(-g_i)^v U$ and $|W| = 3$. By Lemma 3.4.1, $\nu_{-g_i}(W) = 1$. Thus there exist $f_1, f_2 \in \text{supp}(U)$ such that $g_i = f_1 + f_2$. Obviously, $f_1 f_2 \nmid g_1 \dots g_l$. If $f_1 \mid g_1 \dots g_l$ and $f_2 \mid h_1 h_4 h_5$, then $f_2 = g_i + (-f_1)$. Since f_2 is an element of an atom of length 3, we can substitute f_2 by $g_i(-f_1)$ to get an atom of length 4, a contradiction. Therefore, $f_1 f_2 \mid h_1 h_4 h_5$. If $f_1 f_2 = h_4 h_5$, then $g_i = h_6$, a contradiction. By the symmetry of h_4 and h_5 , we only need to consider $f_1 f_2 = h_1 h_4$. Since $(-g_i)h_1 h_4$ is an atom, $g_i h_5(-h_2)(-h_3)(-h_6)$ can only be a product of two atoms which implies that $g_i \in \{h_2, h_3, h_6\}$, a contradiction. \square

We continue with the following two subcases.

CASE 1.1: $\text{ord}(h_4) > 3$.

By **A1**, we have $(-2h_4)^2 \mid U$ and $(-2h_4)^2 \mid V$. Thus, for any $i \in [1, l]$, $g_i \neq -2h_4$ and $-g_i \neq -2h_4$. Therefore, we obtain that $h_1 = h_5 = -2h_4$ and $-h_6 = h_4$, and hence $-h_2 = -h_3 = -2h_4$. Then $-2h_4 = h_1 = h_2 + h_3 = 4h_4$ which implies that $\text{ord}(h_4) = 6$. By **A2**, we obtain that $U = g_1 \dots g_l h_4 (-2h_4)^2$ with $\text{ord}(g_1) = \dots = \text{ord}(g_l) = 2$ and $\text{ord}(h_4) = 6$. Since $\text{D}(G) \geq 5$, we get $l \geq 2$. Then $6 + l - 1 = \exp(G) + l - 1 \leq \text{D}(G) = |U| = l + 3$, a contradiction

CASE 1.2: $\text{ord}(h_4) = 3$.

By **A1**, $h_4^2 \mid U$ and $h_4^2 \mid V$. Thus $h_4^4 \mid h_1 h_4 h_5 (-h_2)(-h_3)(-h_6)$. But $h_4^6 \neq h_1 h_4 h_5 (-h_2)(-h_3)(-h_6)$. Without loss of generality, we can assume that $h_4 = h_5 = -h_3 = -h_6$, $h_1 \neq h_4$, and $-h_2 \neq h_4$.

Suppose $\nu_{h_1}(U) > 1$ and $\nu_{-h_2}(V) > 1$. Then $-h_1 \mid V$ and $h_2 \mid U$. Since $h_1 + (-h_2) + h_4 = 0$, then $(-h_1)h_2 h_4^2$ is an atom of length 4 and divides UV , a contradiction. Thus, by symmetry, we may assume that $\nu_{-h_2}(U) = 1$. Then $h_2 \nmid U$. Gathering the g_i 's, which are equal to h_1 and renumbering if necessary we obtain that $U = g_1 \dots g_s h_1^v h_4^2$ with $s \in [0, l]$, $v = \nu_{h_1}(U)$, $\text{ord}(g_1) = \dots = \text{ord}(g_s) = 2$ (by **A2**), and $\text{ord}(h_4) = 3$.

If $s = 0$, then $\nu_{h_1} = h_4$ and $G = \langle h_1 \rangle$, a contradiction.

Suppose $s \geq 1$. Since $2\sigma(U) = 2\nu_{h_1} + 4h_4 = 0$, we obtain that $2h_4 = 2\nu_{h_1}$ and $6\nu_{h_1} = 0$. Since $(-h_1)^{v-1} h_4^2 \mid V$, we obtain that $(-h_1)^{v-1} h_4^2$ is zero-sum free and, for any $j \in [v+1, 2v]$, we have $j h_1 \in \Sigma(((-h_1)^{v-1} h_4^2))$. This implies that $\text{ord}(h_1) > 2v$. Thus $6v/\text{ord}(h_1) < 3$ which implies that $\text{ord}(h_1) = 3v$ or $6v$. But $\sigma(h_1^v h_4^2) = 3\nu_{h_1} \neq 0$, hence $\text{ord}(h_1) = 6v = \exp(G)$. Therefore,

$$\exp(G) + s - 1 \leq \text{D}(G) = |U| = s + v + 2 \leq 6v + s - 3 \leq \exp(G) + s - 3,$$

a contradiction.

CASE 2: There exists an $A \in \mathcal{A}(G)$ with $A \mid UV$ and $|A| = 4$.

Then UV has a factorization of the form $UV = W_1 \dots W_{\text{D}(G)-1}$ where $W_1, \dots, W_{\text{D}(G)-1} \in \mathcal{A}(G)$, $|W_{\text{D}(G)-1}| = 4$, and all the other W_i have length 2. Conversely, note that for any $A' \in \mathcal{A}(G)$ with $A' \mid UV$ and $|A'| = 4$, $UV(A')^{-1}$ can only be a product of atoms of length 2 because $\text{D}(G) \geq 5$. We start with the following assertion.

A3. If $g \in \text{supp}(U) \cap \text{supp}(V)$, then $\text{ord}(g) = 2$.

Proof of A3. Let $U = g_1 \cdots g_l h_1 h_2$ and $V = (-g_1) \cdots (-g_l) h_3 h_4$ where $l \in \mathbb{N}$, $g_1, \dots, g_l, h_1, \dots, h_4 \in G$ not necessarily distinct and $W_{D(G)-1} = h_1 h_2 h_3 h_4$. Let $g \in \text{supp}(U) \cap \text{supp}(V)$. If $g = g_i$ for some $i \in [1, l]$, then $g | V$, $-g | V$, and hence $g = -g$. If $g = -g_i$ for some $i \in [1, l]$, then $g | U$, $-g | U$, and hence $g = -g$.

It remains to consider the case where $g \in \{h_1, h_2\} \cap \{h_3, h_4\}$, say $h_1 = h_3 = g$. We will show that this is not possible. Assume this holds, choose an $h \in \text{supp}(U) \setminus \{g\}$, and consider the sequence $X = h^{-1} U h_3$. Since $|X| = D(G)$, there exists an atom $A \in \mathcal{A}(G)$ such that $A | X$ and $|A| \in [3, 4]$. Note that $h_1 h_3 | A$.

Suppose that $|A| = 3$. Then there exists $h' \in \text{supp}(U)$ such that $h' = -h_1 - h_3 = -2g$. If $h' = h_2$, then $h_1 h_2 h_3$ is a proper zero-sum sequence of $W_{D(G)-1}$, a contradiction. Otherwise, $h' \in \{g_1, \dots, g_l\}$, hence $-h' = 2g \in \text{supp}(V)$, a contradiction to $h_3 = g \in \text{supp}(V)$ (recall Lemma 3.4).

Suppose that $|A| = 4$. If $h_2 | A$, then (note that $h_1 + h_2 + h_3 + h_4 = 0$) $A = h_1 h_2 h_3 h_4$ and $-h_4 | V$, a contradiction. Thus $h_2 \nmid A$, and the similar argument shows that $h_4 \nmid A$. This implies that $A = g_i g_j h_1 h_3$ with $i, j \in [1, l]$. Then A and $A' = (-g_i)(-g_j)h_2 h_4$ are two atoms of length 4 dividing UV , a contradiction. \square

Clearly, there are precisely two possibilities for U and V which will be discussed in the following two subcases.

CASE 2.1: $U = g_1^{k_1} \cdots g_l^{k_l} h_1^{r_1-1} h_2^{r_2-1} h_3^{r_3-1} h_4^{r_4-1} h_1 h_2$ and

$$V = (-g_1)^{k_1} \cdots (-g_l)^{k_l} (-h_1)^{r_1-1} (-h_2)^{r_2-1} (-h_3)^{r_3-1} (-h_4)^{r_4-1} (-h_3)(-h_4), \text{ where}$$

$k_1, \dots, k_l, r_1, \dots, r_4 \in \mathbb{N}$, $g_1, \dots, g_l, h_1, \dots, h_4$ are pairwise distinct, and $h_1 h_2 (-h_3)(-h_4) = W_{D(G)-1}$.

We start with the following assertions.

A4. For each $g_i \in \{g_1, \dots, g_l\}$ with $\text{ord}(g_i) > 2$, we have that $g_i = h_1 + h_2$.

A5. $r_1 = r_2 = r_3 = r_4 = 1$.

A6. $\text{ord}(g_i) = 2$ for all $i \in [1, l]$.

Proof of A4. Let $i \in [1, l]$ such that $\text{ord}(g_i) > 2$, say $i = 1$. Then, by Lemma 3.4.2, there exists an atom $A \in \mathcal{A}(G)$ such that $A | (-g_1)^{k_1} g_1^{-k_1} U$ and $|A| \in [3, 4]$. We distinguish four subcases depending on the multiplicity of $v_{-g_1}(A)$ and on $|A|$.

Suppose that $v_{-g_1}(A) = 3$. Then $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 | A$ or $h_2 | A$. Then $A = (-g_1)^3 h_i$ with $i \in [1, 2]$, say $i = 1$. It follows that $A' = g_1^3 h_2 (-h_3)(-h_4)$ is a zero-sum subsequence of $UV(A)^{-1}$ which implies that A' is a product of atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 2$. Then $|A| = 4$ by Lemma 3.4.1. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 | A$ or $h_2 | A$. If $h_1 h_2 | A$, then $A = (-g_1)^2 h_1 h_2$, and hence $A' = g_1^2 (-h_3)(-h_4)$ is an atom of length 4 and divides UVA^{-1} , a contradiction. By symmetry, we may assume that $h_1 | A$ and $h_2 \nmid A$. Thus $A = (-g_1)^2 h_1 h$, where $h \in \{g_2, \dots, g_l, h_1, h_3, h_4\}$ and $(-h)(-h_3)(-h_4) | V$. Therefore $A' = g_1^2 h_2 (-h_3)(-h_4)(-h)$ is a zero-sum subsequence of UVA^{-1} which implies that A' is a product of three atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 1$ and $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 | A$ or $h_2 | A$. If $h_1 h_2 | A$, then $A = (-g_1) h_1 h_2 h$, where $h \in \{g_2, \dots, g_l, h_1, h_2, h_3, h_4\}$ and $(-h)(-h_3)(-h_4) | V$. Hence $A' = g_1 (-h_3)(-h_4)(-h)$ is an atom of length 4 and divides UVA^{-1} , a contradiction. By symmetry, we may assume that $h_1 | A$ and $h_2 \nmid A$. Thus $A = (-g_1) h_1 h h'$ where $h, h' \in \{g_2, \dots, g_l, h_1, h_3, h_4\}$ and $(-h)(-h')(-h_3)(-h_4) | V$. Thus $A' = g_1 (-h)(-h') h_2 (-h_3)(-h_4)$ is a zero-sum subsequence of UVA^{-1} which implies that A' is a product of three atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 1$ and $|A| = 3$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 | A$ or $h_2 | A$. If $h_1 h_2 \nmid A$, by symmetry we may assume that $h_1 | A$ and $h_2 \nmid A$. Thus $A = (-g_1) h_1 h$, where $h \in \{g_2, \dots, g_l, h_1, h_3, h_4\}$ and $(-h)(-h_3)(-h_4) | V$. It follows that $A' = g_1 (-h) h_2 (-h_3)(-h_4)$ is a zero-sum subsequence of UVA^{-1} which implies that A' is a product of two atoms, a contradiction. Therefore, $h_1 h_2 | A$ and $g_1 = h_1 + h_2$. \square

Proof of A5. By symmetry it is sufficient to show that $r_3 = 1$. Assume to the contrary that $r_3 \geq 2$. We proceed in several steps.

(i) In the first step we will show that $h_3 = g_i + h_1$ for some $i \in [1, l]$.

By Lemma 3.4.2, there exists an atom $A \in \mathcal{A}(G)$ such that $A \mid h_3^{-(r_3-1)}(-h_3)^{r_3-1}U$ and $|A| \in [3, 4]$. We distinguish four subcases depending on the multiplicity of $v_{-g_1}(A)$ and on $|A|$.

Suppose that $v_{-h_3}(A) = 3$. Then $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 \mid A$ or $h_2 \mid A$. Then $A = (-h_3)^3 h_i$ with $i \in [1, 2]$, say $i = 1$. Therefore $A' = h_3^2 h_2(-h_4)$ is an atom of length 4 and divides UVA^{-1} , a contradiction.

Suppose that $v_{-h_3}(A) = 2$. Then $|A| = 4$ by Lemma 3.4.1. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 \mid A$ or $h_2 \mid A$. If $h_1 h_2 \mid A$, then $A = (-h_3)^2 h_1 h_2$ which implies that $h_3 = h_4$, a contradiction. By symmetry, we may assume that $h_1 \mid A$ and $h_2 \nmid A$. Thus $A = (-h_3)^2 h_1 h$, where $h \in \{g_1, \dots, g_l, h_1, h_4\}$ and $(-h)(-h_3)(-h_4) \mid V$. Therefore $A' = h_3^2 h_2(-h_3)(-h_4)(-h)$ is a zero-sum subsequence of UVA^{-1} which implies that A' is a product of three atoms of length 2, a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 \mid A$ or $h_2 \mid A$. If $h_1 h_2 \mid A$, then $A = (-h_3)h_1 h_2 h$ with $h \in \{g_1, \dots, g_l, h_1, h_2, h_4\}$ and $-h \mid V$. Hence $h = -h_4$ and $-h = h_4 \mid V$, a contradiction. By symmetry, we may assume that $h_1 \mid A$ and $h_2 \nmid A$. Thus $A = (-h_3)h_1 h h'$ where $h, h' \in \{g_1, \dots, g_l, h_1, h_4\}$ and $(-h)(-h')(-h_3)(-h_4) \mid V$. Thus $A' = (-h)(-h')h_2(-h_4)$ is an atom of length 4 and divides UVA^{-1} , a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 3$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h_1 \mid A$ or $h_2 \mid A$. If $h_1 h_2 \mid A$, then $A = (-h_3)h_1 h_2$ which implies that $h_4 = 0$, a contradiction. Thus, by symmetry, we may assume that $h_1 \mid A$ and $h_2 \nmid A$. Then $A = (-h_3)h_1 h$, where $h \in \{g_1, \dots, g_l, h_1, h_4\}$. By Lemma 3.4.1, we obtain that $h \notin \{h_1, h_4\}$. Therefore, $h_3 = h_1 + g_i$ for some $i \in [1, l]$.

(ii) In the second step we will show that $\text{ord}(g_j) = 2$ for all $j \in [1, l]$ and $r_1 = 1$.

Assume to the contrary that there is a $j \in [1, l]$ such that $\text{ord}(g_j) > 2$. Then $g_j = h_1 + h_2$ by **A4** and hence $A_1 = g_j(-h_3)(-h_4)$ and $A_2 = g_i(-h_3)h_1$ are two atoms of length 3 and divide UV . It follows that $UV(A_1 A_2)^{-1}$ is a product of atoms of length 2, but $v_{h_2}(UV(A_1 A_2)^{-1}) = v_{h_2}(UV) > v_{-h_2}(UV) = v_{-h_2}(UV(A_1 A_2)^{-1})$, a contradiction.

Assume to the contrary that $r_1 \geq 2$. Since $\text{ord}(g_i) = 2$ and $h_3 = h_1 + g_i$, we obtain that $h_1^2(-h_3)^2$ and $(-h_1)h_3 g_i$ are two atoms and divide UV , a contradiction.

(iii) In the third step we show that $\text{ord}(h_3) = 4r_3$.

Consider the sequence $X = U(h_3^{r_3-1}h_1)^{-1}(-h_3)^{r_3}$. Since $|X| = D(G)$, there exists an atom $A \in \mathcal{A}(G)$ with $A \mid X$ and $|A| \in \{3, 4, D(G)\}$. We distinguish three subcases depending on $|A|$.

Suppose that $|A| = 3$. Since $A \nmid UV(W_{D(G)-1})^{-1}$ and $r_1 = 1$, we must have that $h_2 \mid A$. Thus $A = (-h_3)h_2 h$, where $h \in \{g_1, \dots, g_l, h_2, -h_3\}$. By Lemma 3.4.1, $h \notin \{h_2, -h_3\}$. Therefore, A and $A' = (-h_3)h_1 g_i$ are two atoms of length 3 and divide UV . It follows that $UV(AA')^{-1}$ is a product of atoms of length 2 but $v_{h_4}(UV(AA')^{-1}) < v_{-h_4}(UV(AA')^{-1})$, a contradiction.

Suppose that $|A| = 4$. Then UVA^{-1} is a product of atoms of length 2, but $r_1 = 1 = v_{h_1}(UVA^{-1}) > v_{-h_1}(UVA^{-1}) = 0$, a contradiction.

Suppose that $|A| = D(G)$. Then $A = X$ and hence $h_1 = -(2k_3 - 1)h_3$. By steps (i) and (ii), we obtain that $g_i = 2r_3 h_3$ and $4r_3 h_3 = 0$. Since Uh_1^{-1} is zero-sum free and for each $j \in [1, 2r_3 - 1]$, $jh_3 \in \Sigma(Uh_1^{-1})$, then $\text{ord}(h_3) > 2r_3$ which implies that $\text{ord}(h_3) = 4r_3$.

(iv) In the final step we will obtain a contradiction to our assumption that $r_3 \geq 2$. Clearly, similar arguments as given in the steps (i), (ii), and (iii) show that $r_2 \geq 2$ implies that $\text{ord}(h_2) = 4r_2$, and that $r_4 \geq 2$ implies that $\text{ord}(h_4) = 4r_4$. We proceed with the following four subcases depending on r_2 and r_4 .

Suppose that $r_2 \geq 2$ and $r_4 \geq 2$. Since $h_3 = h_1 + g_i$ and $\text{ord}(g_i) = 2$, we obtain that $h_2 = h_4 + g_i$ and hence $2h_2 = 2h_4$. Therefore, $(-h_2)h_4 g_i$ and $h_2^2(-h_4)^2$ are two atoms and divide UV , a contradiction.

Suppose that $r_2 = r_4 = 1$. Then $U = g_1 \cdots g_l h_3^{r_3-1} h_1 h_2$ and

$$\exp(G) + l - 1 \leq D(G) = l + r_3 + 1 < l - 1 + 4r_3 \leq \exp(G) + l - 1,$$

a contradiction.

Suppose that $r_2 = 1$ and $r_4 \geq 2$. Then $U = g_1 \cdots g_l h_3^{r_3-1} h_4^{r_4-1} h_1 h_2$ and

$$\exp(G) + l - 1 \leq D(G) = l + r_3 + r_4 < l - 1 + 4 \max\{r_3, r_4\} \leq \exp(G) + l - 1,$$

a contradiction.

Suppose that $r_4 = 1$ and $r_2 \geq 2$. Then $U = g_1 \cdots g_l h_3^{r_3-1} h_2^{r_2-1} h_1 h_2$ and

$$\exp(G) + l - 1 \leq D(G) = l + r_3 + r_2 < l - 1 + 4 \max\{r_2, r_3\} \leq \exp(G) + l - 1,$$

a contradiction. \square

Proof of A6. Assume the contrary that there is an $i \in [1, l]$ such that $\text{ord}(g_i) > 2$, say $i = 1$. Then A4 implies that $g_1 = h_1 + h_2$. Since g_1, \dots, g_l are pairwise distinct, it follows that $\text{ord}(g_2) = \dots = \text{ord}(g_l) = 2$ and $k_2 = \dots = k_l = 1$. Thus $U = g_1^{k_1} g_2 \cdots g_l h_1 h_2$, $0 = \sigma(U) = k_1 g_1 + g_2 + \dots + g_l + g_1$, whence $\text{ord}((k_1 + 1)g_1) = 2$ and $\text{ord}(g_1) = 2(k_1 + 1) \geq 4$. It follows that

$$\exp(G) + l - 1 \leq D(G) = |U| = k_1 + l + 1 \leq \frac{\text{ord}(g_1)}{2} + l \leq \text{ord}(g_1) + l - 2 < \exp(G) + l - 1,$$

a contradiction. \square

Now by A4, A5, and A6, U has the form $U = g_1 \cdots g_l h_1 h_2$ with $\text{ord}(g_i) = 2$ for each $i \in [1, l]$. If $\exp(G) \geq 4$, then

$$\exp(G) + l - 1 \leq D(G) = |U| = l + 2 < 4 + l - 1,$$

a contradiction. Thus G must be an elementary 2-group. Since (g_1, \dots, g_l, h_1) is a basis of G , then $h_2 = g_1 + \dots + g_l + h_1$ and $h_1 + h_2 = h_3 + h_4 = g_1 + \dots + g_l$. We can assume that $h_3 = h_1 + \sum_{i \in I} g_i$ for some $\emptyset \neq I \subsetneq [1, l]$ and $h_4 = h_1 + \sum_{i \in [1, l] \setminus I} g_i$. Then $A_1 = h_3 h_1 \prod_{i \in I} g_i$ and $A_2 = h_1 h_4 \prod_{i \in [1, l] \setminus I} g_i$ are two atoms of lengths $|A_1|, |A_2| \in [3, D(G) - 1]$. If $i \in [1, 2]$ and $|A_i| = 4$, then $UV A_i^{-1}$ has to be a product of atoms of length 2, a contradiction. Thus it follows that $|A_1| = |A_2| = 3$ whence $l = 2$ which implies that $D(G) = 4$, a contradiction.

CASE 2.2: $U = g_1^{k_1} \cdots g_l^{k_l} h^{r-2} h_3^{r_3-1} h_4^{r_4-1} h^2$ and

$$V = (-g_1)^{k_1} \cdots (-g_l)^{k_l} (-h)^{r-2} (-h_3)^{r_3-1} (-h_4)^{r_4-1} (-h_3)(-h_4), \quad \text{where}$$

$k_1, \dots, k_l, r_1, \dots, r_4 \in \mathbb{N}$, $g_1, \dots, g_l, h, h_3, h_4$ are pairwise distinct with the only possible exception that $h_3 = h_4$ may hold, and $h^2(-h_3)(-h_4) = W_{D(G)-1}$.

We start with the following assertions.

A7. For each $i \in [1, l]$ we have $\text{ord}(g_i) = 2$.

A8. $h_3 = h_4$.

Proof of A7. Assume to the contrary that there is an $i \in [1, l]$, say $i = 1$, such that $\text{ord}(g_1) > 2$.

Then, by Lemma 3.4.2, there exists an atom $A \in \mathcal{A}(G)$ such that $A | (-g_1)^{k_1} g_1^{-k_1} U$ and $|A| \in [3, 4]$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. We distinguish four subcases depending on the multiplicity of $v_{-g_1}(A)$ and on $|A|$.

Suppose that $v_{-g_1}(A) = 3$. Then $|A| = 4$ and $A = (-g_1)^3 h$. It follows that $A' = g_1^3 h(-h_3)(-h_4)$ is a zero-sum subsequence of $UV(A)^{-1}$ which implies that A' is a product of atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 2$. Then $|A| = 4$ by Lemma 3.4.1. If $h^2 | A$, then $A = (-g_1)^2 h^2$, and hence $A' = g_1^2(-h_3)(-h_4)$ is an atom of length 4 and divides $UV A^{-1}$, a contradiction. If $h^2 \nmid A$, we obtain that $A = (-g_1)^2 h f$, where $f \in \{g_2, \dots, g_l, h_3, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. Therefore $A' = g_1^2 h(-h_3)(-h_4)(-f)$ is a zero-sum subsequence of $UV A^{-1}$ which implies that A' is a product of three atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 1$ and $|A| = 4$. If $h^2 | A$, then $A = (-g_1) h^2 f$, where $f \in \{g_2, \dots, g_l, h, h_3, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. Hence $A' = g_1(-h_3)(-h_4)(-f)$ is an atom of length 4 and divides $UV A^{-1}$, a contradiction. If $h^2 \nmid A$, then $A = (-g_1) h f f'$ where $f, f' \in \{g_2, \dots, g_l, h_3, h_4\}$ and $(-f)(-f')(-h_3)(-h_4) | V$. Thus $A' = g_1(-f)(-f')h(-h_3)(-h_4)$ is a zero-sum subsequence of $UV A^{-1}$ which implies that A' is a product of three atoms of length 2, a contradiction.

Suppose that $v_{-g_1}(A) = 1$ and $|A| = 3$. If $h^2 | A$, then $g_1 = 2h$, a contradiction to Lemma 3.4.1. If $h^2 \nmid A$, then $A = (-g_1)hf$, where $f \in \{g_2, \dots, g_l, h_3, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. It follows that $A' = g_1(-f)h(-h_3)(-h_4)$ is a zero-sum subsequence of $UV A^{-1}$ which implies that A' is a product of two atoms, a contradiction. \square

Proof of A8. Assume to the contrary that $h_3 \neq h_4$. If $r_3 = r_4 = 1$, then $U = g_1 \cdot \dots \cdot g_l h^r$ with $l \geq 1$, $\text{ord}(h) = 2r \geq 4$, and hence

$$\exp(G) + l - 1 \leq D(G) = |U| = l + r \leq 2r + l - 4 < \exp(G) + l - 1,$$

a contradiction. Thus after renumbering if necessary we may assume that $r_3 \geq 2$. We will show this is impossible.

By Lemma 3.4, there exists an atom $A \in \mathcal{A}(G)$ such that $A | h_3^{-(r_3-1)}(-h_3)^{r_3-1}U$ and $|A| \in [3, 4]$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. We distinguish four subcases depending on the multiplicity of $v_{-h_3}(A)$ and on $|A|$.

Suppose that $v_{-h_3}(A) = 3$. Then $|A| = 4$ and $A = (-h_3)^3 h$. It follows that $A' = h_3^2 h(-h_4)$ is an atom and divides $UV(A)^{-1}$, a contradiction.

Suppose that $v_{-h_3}(A) = 2$. Then $|A| = 4$ by Lemma 3.4.1. If $h^2 | A$, then $A = (-h_3)^2 h^2$ which implies that $h_3 = h_4$, a contradiction. If $h^2 \nmid A$, we obtain that $A = (-h_3)^2 h f$, where $f \in \{g_1, \dots, g_l, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. Therefore $A' = h_3 h(-h_4)(-f)$ is an atom and divides $UV A^{-1}$, a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 4$. If $h^2 | A$, then $A = (-h_3)h^2 f$, where $f \in \{g_1, \dots, g_l, h, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. Hence $h_4 = 2h - h_3 = -f | V$, a contradiction. If $h^2 \nmid A$, then $A = (-h_3)h f f'$ where $f, f' \in \{g_1, \dots, g_l, h_4\}$ and $(-f)(-f')(-h_3)(-h_4) | V$. Thus $A' = (-f)(-f')h(-h_4)$ is an atom and divides $UV A^{-1}$, a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 3$. Since $h^2(-h_3)(-h_4)$ is an atom, we obtain that $h^2 \nmid A$, and hence $A = (-h_3)h f$, where $f \in \{g_1, \dots, g_l, h_4\}$ and $(-f)(-h_3)(-h_4) | V$. If $f = h_4$, then $h = 2h_4$ which implies a contradiction to Lemma 3.4.1 (recall $f = h_4 | U$ and $h = 2h_4 | U$). If $f \neq h_4$, by **A7** $\text{ord}(f) = 2$ and hence $2h = 2h_3$ which implies that $h_3 = h_4$, a contradiction.

Suppose that $v_{-h_3}(A) = 3$. Then $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. Then $h_3^2 h(-h_4)$ is another atom of length 4, a contradiction.

Suppose that $v_{-h_3}(A) = 2$. Then $|A| = 4$ by Lemma 3.4.1. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. If $A = (-h_3)^2 g_i h$, $A' = h_3(-g_i)h(-h_4)$ is zero-sum and divide $UV(A)^{-1}$, then A' is a product of two atoms of length 2, a contradiction. It follows that A can only be $(-h_3)^2 h^2$ which implies that $h_3 = h_4$, a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 4$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. If $h^2 | A$, let $A = (-h_3)h^2 g_i$, then $g_i = -h_4$ which implies that $h_4 = -g_i | V$, a contradiction. So we can assume that $v_h(A) = 1$. Let $A = (-h_3)g_i g_j h$, then $2h = 2h_3$ which implies that $h_3 = h_4$, a contradiction.

Suppose that $v_{-h_3}(A) = 1$ and $|A| = 3$. Since $A \nmid UV(W_{D(G)-1})^{-1}$, we must have that $h | A$. Obviously $A \neq (-h_3)h^2$, so $A = (-h_3)g_i h$ with $i \in [2, l]$. Then $2h = 2h_3$ which implies $h_3 = h_4$, a contradiction. \square

Now, by **A7** and **A8**, U and V have the form

$$U = g_1 \cdot \dots \cdot g_l h_3^{r_3-2} h^r \quad \text{and} \quad V = g_1 \cdot \dots \cdot g_l (-h)^{r-2} (-h_3)^{r_3},$$

where $l \geq 0$, $r_3 \geq 2$, $r \geq 2$, $g_1, \dots, g_l, h_3, h \in G$ are pairwise distinct and $2h = 2h_3$.

If $r_3 = 2$, then $U = g_1 \cdot \dots \cdot g_l h^r$, $\text{ord}(h) = 2r$, and

$$\exp(G) + l - 1 \leq D(G) = |U| = l + r \leq 2r + l - 2,$$

a contradiction. Considering V and assuming $r = 2$ we end again up at a contradiction. Therefore we obtain that $r_3 \geq 3$ and $r \geq 3$.

If $r_3 \geq 4$ and $r \geq 4$, then $h^2(-h_3)^2$ and $(-h)^2 h_3^2$ are two atoms of length 4 and divide UV , a contradiction.

Thus by symmetry, we may assume that $r_3 = 3$ and $r \geq 3$.

Suppose that $l = 0$. Then $\sigma(U) = h_3 + rh = 0$ which implies that $G = \langle h \rangle$, a contradiction.

Suppose that $l \geq 1$. Then $2h_3 + 2rh = 0$ which implies that $2(r+1)h = 0$. Thus $\text{ord}(h) = 2(r+1)$ or $\text{ord}(h) = r+1$. If $\text{ord}(h) = 2(r+1)$, then

$$\exp(G) + l - 1 \leq \text{D}(G) = |U| = l + 1 + r < l - 1 + 2(r+1) \leq \exp(G) + l - 1,$$

a contradiction. If $\text{ord}(h) = r+1$, then $h = h_3 + g_1 + \dots + g_l$ and hence $(-h)h_3g_1 \dots g_l$ is an atom and divides $UV(h^2(-h_3)^2)^{-1}$, a contradiction \square

Proposition 3.7. *Let G be a finite abelian group with $\text{D}(G) \geq 5$. Then the following statements are equivalent:*

- (a) G is either an elementary 2-group, or a cyclic group, or isomorphic to $C_2 \oplus C_{2n}$ with $n \geq 2$.
- (b) There exist $U, V \in \mathcal{A}(G)$ with $\text{L}(UV) = \{2, \text{D}(G) - 1\}$ and $|U| - 1 = |V| = \text{D}(G) - 1$.

Proof. (a) \Rightarrow (b) Suppose that G is an elementary 2-group and let (e_1, \dots, e_r) be a basis of G with $\text{ord}(e_1) = \dots = \text{ord}(e_r) = 2$. Then $\text{D}(G) = r + 1$, $U = e_1 \dots e_r e_0 \in \mathcal{A}(G)$, where $e_0 = e_1 + \dots + e_r$, $V = e_1 \dots e_{r-1}(e_0 + e_r) \in \mathcal{A}(G)$, and $\text{L}(UV) = \{2, r\}$.

Suppose that G is cyclic, and let $e \in G$ with $\text{ord}(e) = |G| = \text{D}(G)$. Then $U = e^{|G|} \in \mathcal{A}(G)$, $V = (-e)^{|G|-1}(-2e) \in \mathcal{A}(G)$, and $\text{L}(UV) = \{2, |G| - 1\}$.

Suppose that G is isomorphic to $C_2 \oplus C_{2n}$ with $n \geq 2$, and let (e_1, e_2) be a basis of G with $\text{ord}(e_1) = 2$ and $\text{ord}(e_2) = 2n$. Then $\text{D}(G) = 2n + 1$, $U = e_1 e_2^{2n-1}(e_1 + e_2) \in \mathcal{A}(G)$, $V = (-e_2)^{2n} \in \mathcal{A}(G)$, and $\text{L}(UV) = \{2, 2n\}$.

(b) \Rightarrow (a) Assume to the contrary that G is neither an elementary 2-group, nor a cyclic group, nor isomorphic to $C_2 \oplus C_{2n}$ for any $n \geq 2$. Then $\text{D}(G) > \exp(G) + 1$.

Let $A \in \mathcal{A}(G)$ with $A | UV$. Then we have $|A| \in \{2, 3, \text{D}(G) - 1, \text{D}(G)\}$. Furthermore, if $|A| = 3$, then UVA^{-1} is a product of atoms of length 2, and if $|A| \in [\text{D}(G) - 1, \text{D}(G)]$, then $UVA^{-1} \in \mathcal{A}(G)$.

Since $\text{L}(UV) = \{2, \text{D}(G) - 1\}$ and $|U| - 1 = |V| = \text{D}(G) - 1$, U and V have the form

$$U = g_1^{k_1} \dots g_s^{k_s} \quad \text{and} \quad V = (-g_1 - g_2)(-g_1)^{k_1-1}(-g_2)^{k_2-1}(-g_3)^{k_3} \dots (-g_s)^{k_s},$$

where $s, k_1, \dots, k_s \in \mathbb{N}$, $g_1, \dots, g_s \in G$ are pairwise distinct with the only possible exception that $g_1 = g_2$ may hold.

Since G is not cyclic, we have $s \geq 2$. Suppose that $s = 2$. If $g_1 = g_2$, then $\text{D}(G) = k_1 + k_2 = \text{ord}(g_1)$ which implies that G is a cyclic group, a contradiction. Suppose that $g_1 \neq g_2$. Since $\text{D}(G) > \exp(G) + 1$, Lemma 3.2.2 implies that $v_{g_1}(U) + v_{-g_1}(V) = k_1 + k_1 - 1 \leq \text{ord}(g_1)$ and $v_{g_2}(U) + v_{-g_2}(V) = k_2 + k_2 - 1 \leq \text{ord}(g_2)$. Then

$$\text{D}(G) = k_1 + k_2 \leq \frac{\text{ord}(g_1) + 1}{2} + \frac{\text{ord}(g_2) + 1}{2} \leq 1 + \exp(G) < \text{D}(G),$$

a contradiction.

Thus from now on we suppose that $s \geq 3$, and we continue with the following assertion.

A1. There exist atoms $U', V' \in \mathcal{A}(G)$ such that $UV = U'V'$, say

$$U' = g_1^{k'_1} \dots g_{s'}^{k'_{s'}} \quad \text{and} \quad V' = (-g'_1 - g'_2)(-g'_1)^{k'_1-1}(-g'_2)^{k'_2-1}(-g'_3)^{k'_3} \dots (-g'_{s'})^{k'_{s'}},$$

where $s', k'_1, \dots, k'_{s'} \in \mathbb{N}$, $g'_1, \dots, g'_{s'} \in G$ are pairwise distinct with the only possible exception that $g'_1 = g'_2$ may hold, and $g'_1 + g'_2 \notin \text{supp}(U')$.

Proof of A1. If $g_1 + g_2 \notin \text{supp}(U)$, then we can choose $U' = U$ and $V' = V$.

Suppose $g_1 + g_2 \in \text{supp}(U)$, say $g_3 = g_1 + g_2$. Then $v_{-g_3}(V) = k_3 + 1 \geq 2$ and hence $\text{ord}(g_3) > 2$. By Lemma 3.4.1 and $|U| = \text{D}(G)$, it follows that $g_1 \neq g_2$. We claim that $k_1 = 1$ or $k_2 = 1$. Indeed, if $k_1 \geq 2$ and $k_2 \geq 2$, then $A = g_1 g_2 (-g_3)$ is an atom and $A^2 | UV$, a contradiction.

Suppose $s = 3$. Without loss of generality, we can assume that $k_1 = 1$. Since $D(G) > \exp(G) + 1$, Lemma 3.2.2 implies that $v_{g_2}(U) + v_{-g_2}(V) = k_2 + k_2 - 1 \leq \text{ord}(g_2)$ and $v_{g_3}(U) + v_{-g_3}(V) = k_3 + k_3 + 1 \leq \text{ord}(g_3)$. Then

$$D(G) = |U| = 1 + k_2 + k_3 \leq 1 + \frac{\text{ord}(g_2) + 1}{2} + \frac{\text{ord}(g_3) - 1}{2} \leq 1 + \exp(G) < D(G),$$

a contradiction.

Thus we obtain that $s \geq 4$. Since $g_3^{-1}U$ is a zero-sum free sequence of length $D(G) - 1$, there exists a subsequence T_1 of $g_3^{-1}U$ such that $\sigma(T_1) = (k_3 + 1)g_3$. Therefore, $(-g_3)^{k_3+1}T_1$ is a zero-sum sequence. Thus T_1 has the form $T_1 = g_3^t T_2$ with $t \in [0, k_3 - 1]$ and $T_2 | g_1^{k_1} g_2^{k_2} g_4^{k_4} \cdots g_s^{k_s}$. Then $(-g_3)^{k_3+1-t}T_2$ is a zero-sum sequence without zero-sum subsequences of length 2 which implies that $(-g_3)^{k_3+1-t}T_2$ is an atom of length $|(-g_3)^{k_3+1-t}T_2| \in \{3, D(G) - 1, D(G)\}$. Since $k_3 + 1 - t \geq 2$, $(-g_3)^{k_3+1-t}T_2$ cannot be an atom of length 3 by Lemma 3.4.1, hence $t = 0$, $T_1 = T_2$, and $(-g_3)^{k_3+1}T_1$ can only be an atom of length $D(G) - 1$ or $D(G)$. It follows that $((-g_3)^{k_3+1}T_1)^{-1}UV$ is also an atom and hence $g_4^{k_4} \cdots g_s^{k_s} | T_1$.

Thus any sequence T with $T | g_3^{-1}U$ and $\sigma(T) = (k_3 + 1)g_3$ has the property that $g_4^{k_4} \cdots g_s^{k_s} | T$.

We continue with the following two subcases depending on $|(-g_3)^{k_3+1}T_1|$.

Suppose $|(-g_3)^{k_3+1}T_1| = D(G) - 1$. Since $((-g_3)^{k_3+1}T_1)^{-1}UV$ is an atom, we obtain that $k_1 = k_2 = 1$ and $T_1 = (g_1 g_2 g_3^{k_3})^{-1}U$ which implies that $2(k_3 + 1)g_3 = 0$. Since $g_4^{-1}U$ is a zero-sum free sequence of length $D(G) - 1$, there exists a subsequence W_1 of $g_4^{-1}U$ such that $\sigma(W_1) = (k_3 + 2)g_3$. If $g_3 \nmid W_1$, then $g_3^{k_3}W_1$ is a proper zero-sum subsequence of U , a contradiction. If $g_3 | W_1$, then $g_3^{-1}W_1 | g_3^{-1}U$ and $\sigma(g_3^{-1}W_1) = (k_3 + 1)g_3$, but $g_4^{k_4} \cdots g_s^{k_s} \nmid g_3^{-1}W_1$, a contradiction.

Suppose $|(-g_3)^{k_3+1}T_1| = D(G)$. Since $((-g_3)^{k_3+1}T_1)^{-1}UV$ is an atom, we obtain that $(k_1 = 1$ and $T_1 = (g_1 g_3^{k_3})^{-1}U$) or $(k_2 = 1$ and $T_1 = (g_2 g_3^{k_3})^{-1}U$). By symmetry, we may assume that $k_1 = 1$, $T_1 = (g_1 g_3^{k_3})^{-1}U$, and hence $g_1 = (-2k_3 - 1)g_3$. Choose

$$\begin{aligned} U' &= (-g_3)^{k_3+1}T_1 = g_2(-g_3)g_2^{k_2-1}(-g_3)^{k_3}g_4^{k_4} \cdots g_s^{k_s} \quad \text{and} \\ V' &= ((-g_3)^{k_3+1}T_1)^{-1}UV = g_1(-g_2)^{k_2-1}g_3^{k_3}(-g_4)^{k_4} \cdots (-g_s)^{k_s}, \end{aligned}$$

then U', V' are two atoms with $U'V' = UV$ and $g_2 + (-g_3) = -g_1 \notin \text{supp}(U')$. \square

Thus from now on we may assume that $g_1 + g_2 \notin \text{supp}(U)$, and recall that $s \geq 3$. We continue with three further assertions.

A2. $(\text{supp}(U) + \text{supp}(U)) \cap (\text{supp}(U) \setminus \{g_1, g_2\}) = \emptyset$.

A3. Let $i \in [3, s]$ with $\text{ord}(g_i) > 2$. Then $(-2k_i g_i = g_1$ and $k_1 = 1)$ or $(-2k_i g_i = g_2$ and $k_2 = 1)$.

A4. If $k_1 = 1$, then $(g_2 = -2g_1$ and $k_2 = 1)$ or $(2g_1 + 2g_2 = 0$ and $k_2 = 1)$. If $k_2 = 1$, then $(g_1 = -2g_2$ and $k_1 = 1)$ or $(2g_1 + 2g_2 = 0$ and $k_1 = 1)$.

Proof of A2. Assume to the contrary that there exists an element $h \in (\text{supp}(U) + \text{supp}(U)) \cap (\text{supp}(U) \setminus \{g_1, g_2\})$. Thus there exist $i, j \in [1, s]$ such that $g_i + g_j = h$ with $h \in \{g_3, \dots, g_s\}$. Lemma 3.4.1 implies that $g_i \neq g_j$. Since $A = (-h)g_i g_j$ is an atom of length 3, then $A^{-1}UV$ is a product of atoms of length 2. It follows that $h = g_1 + g_2 \in \text{supp}(U)$, a contradiction. \square

Proof of A3. By Lemma 3.4.2, there is an $A \in \mathcal{A}(G)$ with $A | g_i^{-k_i}(-g_i)^{k_i}U$ and $|A| \in \{3, D(G) - 1\}$. By **A2** and Lemma 3.4.1, we obtain that $|A| \neq 3$. Thus $|A| = D(G)$. If $A = g_i^{-k_i}(-g_i)^{k_i-1}U$, then $(2k_i - 1)g_i = 0$ which implies that $v_{g_i}(U) + v_{-g_i}(V) = 2k_i > \text{ord}(g_i)$. Lemma 3.2.2 implies a contradiction to $D(G) > \exp(G) + 1$. Hence $A = (-g_i)^{k_i}T$ with $T = g_i^{-k_i}h^{-1}U$, where $h \in \text{supp}(U)$. Since $T^{-1}UV$ is an atom, we obtain that $(k_1 = 1$ and $h = g_1 = -2k_i g_i)$ or $(k_2 = 1$ and $h = g_2 = -2k_i g_i)$. \square

Proof of A4. Suppose that $k_1 = 1$. Then $Y = (-g_1 - g_2)g_2^{k_2} \cdots g_s^{k_s}$ has length $D(G)$. By our assumption on $\text{supp}(U)$, Y has no zero-sum subsequence of length 2. Therefore, Y has a zero-sum subsequence T of length $|T| \in \{3, D(G) - 1, D(G)\}$. We continue with the following three subcases depending on $|T|$. If $|T| = 3$, say $T = (-g_1 - g_2)g_i g_j$ with $i, j \in [2, s]$ not necessarily distinct, then T and $T' = g_1 g_2 (-g_i)(-g_j)$ are two atoms and $T'T | UV$, a contradiction. If $|T| = D(G)$, then $Y = T$ is an atom which implies that

$g_2 = -2g_1$, $-g_1 - g_2 = g_1$, and hence $k_1 = 1$. If $|T| = D(G) - 1$, then $k_2 = 1$, $T = Yg_2^{-1}$, and hence $2g_1 + 2g_2 = 0$.

If $k_2 = 1$, then the assertion follows along the same lines. \square

The remainder of the proof will be divided into the following three cases.

CASE 1: $|\{i \in [3, s] \mid \text{ord}(g_i) > 2\}| \geq 2$, say $\text{ord}(g_3) > 2$ and $\text{ord}(g_4) > 2$.

By **A3**, we can assume that $(k_1 = 1$ and $g_1 = -2k_3g_3 = -2k_4g_4)$ or $(g_1 = -2k_3g_3, g_2 = -2k_4g_4$ and $k_1 = k_2 = 1)$.

Consider the sequence $W = g_1^{k_1}g_2^{k_2}(-g_3)^{k_3}(-g_4)^{k_4}g_5^{k_5} \cdots g_s^{k_s}$. Since $|W| = D(G)$, there exists an atom $Z \in \mathcal{A}(G)$ such that $Z \mid W$ and $|Z| \in \{3, D(G) - 1, D(G)\}$. We distinguish three subcases depending on $|Z|$.

Suppose $|Z| = 3$. By **A2** and Lemma 3.4.1, $g_1 = g_3 + g_4$ or $g_2 = g_3 + g_4$. If $g_1 = g_3 + g_4$, then $(-g_1 - g_2)g_3g_4g_2$ and $g_1(-g_3)(-g_4)$ are two atoms and divide UV , a contradiction. If $g_2 = g_3 + g_4$, then $(-g_1 - g_2)g_1g_3g_4$ and $g_2(-g_3)(-g_4)$ are two atoms and divide UV , a contradiction.

Suppose $|Z| = D(G) - 1$. Since UVZ^{-1} is a atom, we obtain that $Z = Wg_1^{-1}$ or $Z = Wg_2^{-1}$. Therefore, $-2k_3g_3 - 2k_4g_4 = g_1$ or g_2 . Our assumption infers that $-2k_3g_3 - 2k_4g_4 = g_2$ and $g_1 = -2k_3g_3 = -2k_4g_4$ which implies that $2g_1 = g_2$, a contradiction to Lemma 3.4.1.

Suppose $|Z| = D(G)$. Then we obtain that $2k_3g_3 + 2k_4g_4 = 0$. Our assumption infers that $k_1 = 1$, $g_1 = -2k_3g_3 = -2k_4g_4$, and hence $\text{ord}(g_1) = 2$. Therefore, $4k_3g_3 = 0$, $g_1 = 2k_3g_3$, and hence $k_3 \geq 2$ by Lemma 3.4.1. Since $g_1^{-1}U$ is a zero-sum sequence of length $D(G) - 1$, there exists a subsequence W of $g_1^{-1}U$ such that $\sigma(W) = (2k_3 + 1)g_3$. If $g_3 \mid W$, then $g_1g_3^{-1}W$ is a proper zero-sum subsequence of U , a contradiction. Suppose $g_3 \nmid W$. Then $g_1(-g_3)W$ is an atom and divides UV which implies that $|g_1(-g_3)W| \in \{3, D(G) - 1, D(G)\}$. Since $g_3(-g_3) \mid UV(g_1(-g_3)W)^{-1}$, then $UV(g_1(-g_3)W)^{-1}$ is not an atom. Thus $|g_1(-g_3)W| = 3$ which implies that $g_3 \in \text{supp}(U) \cap \text{supp}(V)$, a contradiction to **A2**.

CASE 2: $|\{i \in [3, s] \mid \text{ord}(g_i) > 2\}| = 1$, say $\text{ord}(g_3) > 2$.

By **A3**, we may assume that $k_1 = 1$ and $g_1 = -2k_3g_3$. By **A4**, we obtain that $(g_2 = -2g_1$ and $k_2 = 1)$ or $(2g_1 + 2g_2 = 0$ and $k_2 = 1)$. We continue with the following two subcases.

Suppose that $2g_1 + 2g_2 = 0$ and $k_2 = 1$. Since $\sigma(U) = g_1 + g_2 + k_3g_3 + g_4 + \cdots + g_s = 0$, we obtain that $2k_3g_3 = 0 = -g_1$, a contradiction.

Suppose that $g_2 = -2g_1 = 4k_3g_3$ and $k_2 = 1$. If $s = 3$, then $G = \langle g_3 \rangle$ is cyclic, a contradiction. Hence $s \geq 4$ and $G = \langle g_1, \dots, g_{s-1} \rangle = \langle g_3, \dots, g_{s-1} \rangle$ which implies that $r(G) = s - 3$ and $\exp(G) = \text{ord}(g_3)$ is even. Since $D(G) > \exp(G) + 1$, by Lemma 3.2.2, we infer that $v_{g_3}(U) + v_{-g_3}(V) = 2k_3 \leq \text{ord}(g_3)$. Therefore, $g_1 = -2k_3g_3 \neq 0$ infers that $\text{ord}(g_3) \geq 2k_3 + 2 \geq 4$. Thus

$$\exp(G) + s - 4 \leq D(G) = |U| = k_3 + s - 1 \leq \frac{\text{ord}(g_3)}{2} - 1 + s - 1 \leq \text{ord}(g_3) + s - 4$$

which implies that $\text{ord}(g_3) = 4$, a contradiction to $g_2 = 4k_3g_3 \neq 0$.

CASE 3: $|\{i \in [3, s] \mid \text{ord}(g_i) > 2\}| = 0$.

Since $\sigma(U) = k_1g_1 + k_2g_2 + g_3 + \cdots + g_s = 0$, we obtain that $2k_1g_1 + 2k_2g_2 = 0$.

Suppose that $g_1 = g_2$. Then $\text{ord}(g_1) = 2(k_1 + k_2) \geq 4$. It follows that

$$D(G) = k_1 + k_2 + s - 2 = \frac{\text{ord}(g_1)}{2} + s - 2 < \text{ord}(g_1) - 1 + s - 2 \leq D(G),$$

a contradiction.

Thus $g_1 \neq g_2$. Consider the sequence $S = (-g_1 - g_2)(-g_1)^{k_1-1}g_2^{k_2}g_3 \cdots g_s$. Since $|S| = D(G)$, there exists an atom $Z \in \mathcal{A}(G)$ such that $Z \mid S$ and $|Z| \in \{3, D(G) - 1, D(G)\}$. We distinguish three subcases depending on $|Z|$.

Suppose that $|Z| = D(G)$. Then $Z = S$ and $g_2 = -2k_1g_1 = 2k_2g_2$. Hence $(2k_2 - 1)g_2 = 0$ and $\text{ord}(g_2) = 2k_2 - 1$. If $s = 3$, then $G = \langle g_1 \rangle$ is cyclic, a contradiction. Hence $s \geq 4$ and $G = \langle g_1, \dots, g_{s-1} \rangle = \langle g_1, g_3, \dots, g_{s-1} \rangle$ which implies that $r(G) = s - 2$ and $\text{ord}(g_1) = \exp(G)$ is even. Then $\text{ord}(g_1) \geq 2 \text{ord}(g_2) \geq$

6. Since $D(G) > \exp(G) + 1$, by Lemma 3.2.2, we infer that $v_{g_1}(U) + v_{-g_1}(V) = 2k_1 - 1 \leq \text{ord}(g_1)$ which implies that $2k_1 \leq \text{ord}(g_1)$. Since $g_2 = -2k_1g_1 \neq 0$, we obtain that $2k_1 \leq \text{ord}(g_1) - 2$. Thus

$$\begin{aligned} \exp(G) + s - 3 \leq D(G) = |U| = k_1 + k_2 + s - 2 &\leq \frac{\text{ord}(g_1)}{2} - 1 + \lfloor \frac{\text{ord}(g_1) + 2}{4} \rfloor + s - 2 \\ &\leq \frac{\text{ord}(g_1)}{2} - 1 + \frac{\text{ord}(g_1)}{2} - 1 + s - 2 \leq \text{ord}(g_1) + s - 4 \end{aligned}$$

a contradiction.

Suppose that $|Z| = D(G) - 1$. Then there exists an element $h \mid S$ such that $Z = h^{-1}S$. Since $Z^{-1}UV$ is an atom, we obtain that $(h = -g_1 - g_2)$ or $(h = g_2 \text{ and } k_2 = 1)$. If $h = -g_1 - g_2$, then $Z^{-1}UV = (-g_1 - g_2)g_1^{k_1}(-g_2)^{k_2-1}g_3 \cdots g_s$ is an atom of length $D(G)$ and hence the similar argument of the previous subcase $|Z| = D(G)$ implies a contradiction. Suppose that $h = g_2$ and $k_2 = 1$. By **A4**, we obtain that $k_1 = k_2 = 1$. Thus $\exp(G) + s - 3 \leq D(G) = |U| = s$ which implies that $\exp(G) \leq 3$, a contradiction.

Suppose that $|Z| = 3$. Then UVZ^{-1} can only be a product of atoms of length 2. But $v_{g_1}(UVZ^{-1}) = k_1 > v_{-g_1}(UV) \geq v_{-g_1}(UVZ^{-1})$, a contradiction. \square

Proposition 3.8. *Let G be a finite abelian group with $D(G) \geq 5$. Then the following statements are equivalent:*

- (a) G is either an elementary 2-group, or isomorphic to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$, or isomorphic to $C_2 \oplus C_{2n}$ for some $n \geq 2$.
- (b) There exist $U, V \in \mathcal{A}(G)$ with $L(UV) = \{2, D(G) - 1\}$ and $|U| = |V| = D(G) - 1$.

Proof. (a) \Rightarrow (b) Suppose that G is an elementary 2-group, and let (e_1, \dots, e_r) be a basis of G with $\text{ord}(e_1) = \dots = \text{ord}(e_r) = 2$. Then $D(G) = r + 1$, $U = e_1 \cdots e_{r-1}e_0 \in \mathcal{A}(G)$, where $e_0 = e_1 + \dots + e_{r-1}$, and $L(U(-U)) = \{2, r\}$.

Suppose that G is isomorphic to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$, and let (e_1, \dots, e_r) be a basis of G with $\text{ord}(e_1) = \dots = \text{ord}(e_{r-1}) = 2$ and $\text{ord}(e_r) = 4$. Then $D(G) = r + 3$, $U = e_1 \cdots e_{r-1}e_r^2(e_0 + e_r) \in \mathcal{A}(G)$, where $e_0 = e_1 + \dots + e_r$, and $L(U(-U)) = \{2, r + 2\}$.

Suppose that G is isomorphic to $C_2 \oplus C_{2n}$ for some $n \geq 2$, and let (e_1, e_2) be a basis of G with $\text{ord}(e_1) = 2$ and $\text{ord}(e_2) = 2n$. Then $D(G) = 2n + 1$, $U = e_2^{2n} \in \mathcal{A}(G)$, and $L(U(-U)) = \{2, 2n\}$.

(b) \Rightarrow (a) Clearly, we have $V = -U$, and for every zero-sum sequence W with $W \mid UV$ and $W \neq UV$, it follows that W is either an atom of length $D(G) - 1$ or W is a product of atoms of length 2. We set $U = g_1^{k_1} \cdots g_l^{k_l}$ with $l, k_1, \dots, k_l \in \mathbb{N}$ and $g_1, \dots, g_l \in G$ pairwise distinct.

Suppose that $l = 1$. Then $k_1 = \text{ord}(g_1) = D(G) - 1$. Thus k_1 is even and $G \cong C_2 \oplus C_{k_1}$.

Suppose that $l \geq 2$. For each $i \in [1, l]$, the sequence $S_i = g_i^{-k_i}(-g_i)^{k_i}U$ is either zero-sum free or an atom; clearly, S_i is an atom if and only if $2k_i g_i = 0$. So we can distinguish two cases.

CASE 1: For each $i \in [1, l]$ we have $2k_i g_i = 0$.

We claim that for any $i \in [1, l]$, the tuple $(g_1, \dots, g_{i-1}, g_{i+1}, \dots, g_l)$ is independent. Clearly, it is sufficient to prove the claim for $i = l$. Assume to the contrary that (g_1, \dots, g_{l-1}) is not independent. Then there is an atom W with $W \mid g_1^{k_1} \cdots g_{l-1}^{k_{l-1}} \cdot (-g_1)^{k_1} \cdots (-g_{l-1})^{k_{l-1}}$ and $|W| > 2$. Then W is an atom of length $D(G) - 1$, and thus $W^{-1}UV$ is also an atom of length $D(G) - 1$. But $g_l(-g_l) \mid W^{-1}UV$, a contradiction. After renumbering if necessary we may suppose that

$$\text{ord}(g_l) = \min\{\text{ord}(g_1), \dots, \text{ord}(g_l)\} \quad \text{and} \quad \text{ord}(g_1) = \min\{\text{ord}(g_1), \dots, \text{ord}(g_{l-1})\}.$$

Suppose that $l = 2$. By our assumption that $\text{ord}(g_1) \geq \text{ord}(g_2)$, we obtain that

$$\begin{aligned} \exp(G) - 1 \leq D(G) - 1 = k_1 + k_2 &= \frac{\text{ord}(g_1)}{2} + \frac{\text{ord}(g_2)}{2} = \text{ord}(g_1) - \frac{\text{ord}(g_1) - \text{ord}(g_2)}{2} \\ &\leq \exp(G) - \frac{\text{ord}(g_1) - \text{ord}(g_2)}{2} \leq \exp(G). \end{aligned}$$

If $D(G) = \exp(G)$, then $\text{ord}(g_1) = \exp(G)$, $\text{ord}(g_1) - \text{ord}(g_2) = 2$, and hence $\text{ord}(g_2) | 2$ which implies $\text{ord}(g_2) = 2$ and $\text{ord}(g_1) = 4 = D(G) = \exp(G)$, a contradiction to $D(G) \geq 5$. If $D(G) = \exp(G) + 1$, then $G \cong C_2 \oplus C_{2n}$ for some $n \geq 2$.

Suppose that $l \geq 3$. Then

$$\begin{aligned} D(G) - 1 = |U| = k_1 + \dots + k_l &= \frac{\text{ord}(g_1)}{2} + \dots + \frac{\text{ord}(g_l)}{2} \leq \text{ord}(g_1) + \frac{\text{ord}(g_2)}{2} + \dots + \frac{\text{ord}(g_{l-1})}{2} \\ &\leq \text{ord}(g_1) + \dots + \text{ord}(g_{l-1}) - (l-2) \leq D(G). \end{aligned}$$

Suppose that equality holds at the second inequality sign. Then

$$\text{ord}(g_1) + \frac{\text{ord}(g_2)}{2} + \dots + \frac{\text{ord}(g_{l-1})}{2} = \text{ord}(g_1) + \dots + \text{ord}(g_{l-1}) - (l-2).$$

Since $\text{ord}(g_i)/2 \leq \text{ord}(g_i) - 1$ for all $i \in [2, l-1]$, it follows that then $\text{ord}(g_i) = 2$ for all $i \in [2, l-1]$. Because our assumption on the order of the elements, we infer that $\text{ord}(g_1) = \text{ord}(g_l) = 2$, and hence G is an elementary 2-group.

Suppose that inequality holds at the second inequality sign. Then we have

$$\text{ord}(g_1) + \frac{\text{ord}(g_2)}{2} + \dots + \frac{\text{ord}(g_{l-1})}{2} = \text{ord}(g_1) + \dots + \text{ord}(g_{l-1}) - (l-2) - 1.$$

Since $\text{ord}(g_i)/2 \leq \text{ord}(g_i) - 1$ for all $i \in [2, l-1]$, there exists an $i \in [2, l-1]$, say $i = 2$, such that $\text{ord}(g_2) = 4$ and $\text{ord}(g_i) = 2$ for all $i \in [3, l-1]$. If $l = 3$, then $G \cong C_4 \oplus C_4$ or $G \cong C_2 \oplus C_4$, but the first case is a contradiction to Lemma 3.3. If $l \geq 4$, then $G \cong C_2^{l-2} \oplus C_4$.

CASE 2: There exists an $i \in [1, l]$ such that the sequence S_i is zero-sum free, say $i = 1$.

We start with a list of assertions.

- A1.** Let $T, T' \in \mathcal{F}(G)$ be distinct such that $T | U$, $T' | U$, and $\sigma(T) = \sigma(T')$. Then $\text{supp}(T) \cap \text{supp}(T') = \emptyset$, $TT' = U$, and $2\sigma(T) = 0$.
- A2.** Let $T, T' \in \mathcal{F}(G)$ be distinct such that $T | S_1$, $T' | S_1$, and $\sigma(T) = \sigma(T')$. Then $\text{supp}(T) \cap \text{supp}(T') = \emptyset$, $TT' = S_1$, and $2\sigma(T) = -2k_1g_1$.
- A3.** $\Sigma(U) = G$.
- A4.** If $i \in [1, l]$ with $\text{ord}(g_i) > 2$, then $\text{ord}(g_i) > 2k_i$.

Proof of A1. Obviously, $T(-T')$ has sum zero and $T(-T') | UV$ but $T(-T') \neq UV$. So $T(-T')$ must be an atom of length $D(G) - 1$ which implies that $\text{supp}(T) \cap \text{supp}(T') = \emptyset$, $TT' = U$, and $2\sigma(T) = 0$. \square

Proof of A2. Obviously, $T(-T')$ has sum zero, $T(-T') | UV$, and $T(-T') \neq UV$. So $T(-T')$ must be an atom of length $D(G) - 1$ which implies that $\text{supp}(T) \cap \text{supp}(T') = \emptyset$, $TT' = S_1$, and $2\sigma(T) = -2k_1g_1$. \square

Proof of A3. We will show that $|\Sigma(U)| = |G|$. Clearly, we have

$$\begin{aligned} |\Sigma(U)| &= |\{\sigma(T) \mid 1 \neq T \in \mathcal{F}(G), T | U\}| \\ &= |\{g \in G \mid \text{there exist } 1 \neq T \text{ with } T | U \text{ and } \sigma(T) = g\}| \end{aligned}$$

Since $U = g_1^{k_1} \dots g_l^{k_l}$, we have

$$|\{T \in \mathcal{F}(G) \mid 1 \neq T, T | U\}| = (k_1 + 1) \dots (k_l + 1) - 1.$$

By **A1**, there are at most two distinct subsequences of U with given sum $g \in G$. Therefore we obtain

$$\begin{aligned} |\Sigma(U)| &= |\{\sigma(T) \mid 1 \neq T \in \mathcal{F}(G), T | U\}| \\ &= |\{T \in \mathcal{F}(G) \mid 1 \neq T, T | U\}| - \\ &\quad \frac{1}{2} |\{T \in \mathcal{F}(G) \mid T | U \text{ and there is a divisor } T' \text{ of } U \text{ with } T \neq T' \text{ and } \sigma(T) = \sigma(T')\}| \\ &= (k_1 + 1) \dots (k_l + 1) - 1 - |\{T \in \mathcal{F}(G) \mid g_1 \nmid T, T = \prod_{g \in \text{supp}(T)} g^{v_g(U)}, \text{ and } \text{ord}(\sigma(T)) = 2\}|. \end{aligned}$$

Next we study $|\Sigma(S_1)|$. Since S_1 is zero-sum free of length $|S_1| = D(G) - 1$, it follows that $\Sigma(S_1) = G \setminus \{0\}$. Using **A2** for the second equality sign we obtain that

$$\begin{aligned} |G| - 1 &= |\Sigma(S_1)| = (k_1 + 1) \cdots (k_l + 1) - 1 \\ &\quad - |\{T \in \mathcal{F}(G) \mid T = (-g_1)^{k_1} \prod_{g \in \text{supp}(T) \setminus \{-g_1\}} g^{v_g(U)} \text{ and } 2\sigma(T) = -2k_1g_1\}| \\ &= (k_1 + 1) \cdots (k_l + 1) - 1 \\ &\quad - (|\{T \in \mathcal{F}(G) \mid g_1 \nmid T \text{ and } T = \prod_{g \in \text{supp}(T)} g^{v_g(U)}, \text{ ord}(\sigma(T)) = 2\}| + |\{T \in \mathcal{F}(G) \mid T = 1\}|) \\ &= |\Sigma(U)| - 1, \end{aligned}$$

and hence $\Sigma(U) = G$. \square

Proof of A4. Since S_1 is zero-sum free, it follows that $\text{ord}(g_1) > 2k_1$. Now let $i \in [2, l]$ be given and assume to the contrary that $\text{ord}(g_i) \leq 2k_i$. Recall that $\text{ord}(g_i) > v_{g_i}(U) = k_i$. We set $U = g_i^{k_i}U'$. Then $W' = (-g_i)^{\text{ord}(g_i) - k_i}U'$ has sum zero and divides UV . Thus W' is an atom of length $D(G) - 1 = |W'| = |U|$ and hence $\text{ord}(g_i) = 2k_i$. Since $\Sigma(S_1) = G \setminus \{0\}$, S_1 has a subsequence T such that $\sigma(T) = (k_i + 1)g_i$. If $g_i \nmid T$, then $Tg_i^{k_i - 1}$ is a proper zero-sum subsequence of S_1 , a contradiction. If $g_i \mid T$, then $\sigma(g_i^{-1}T) = k_i g_i = \sigma(g_i^{k_i})$. By **A2**, it follows that $0 = 2k_i g_i = 2\sigma(g_i^{-1}T) = -2k_1 g_1 \neq 0$, a contradiction. \square

Now we distinguish two subcases.

CASE 2.1: $|\{i \in [1, l] \mid \text{ord}(g_i) > 2\}| \geq 3$, say $\text{ord}(g_1) > 2$, $\text{ord}(g_2) > 2$, and $\text{ord}(g_3) > 2$.

We start with the following assertion.

A5. There is a subsequence W of U with $\sigma(W) = g_1 - g_2$ such that $W = g_2^{k_2}W'$ and for any $h \mid W'$, $\text{ord}(h) = 2$.

Proof of A5. By **A3**, there exists some $W \in \mathcal{F}(G)$ such that $W \mid U$ and $\sigma(W) = g_1 - g_2$. We claim that $g_1 \nmid W$ but $g_2^{k_2} \mid W$. Assume to the contrary that $g_1 \mid W$. Then $\sigma(g_1^{-1}W) = -g_2 = \sigma(g_2^{-1}U)$. By **A1**, this implies that $2g_2 = 0$, a contradiction. Assume to the contrary that $g_2^{k_2} \nmid W$. Then $g_2W \mid U$ and $\sigma(g_2W) = g_1 = \sigma(g_1)$. By **A1**, this implies that $2g_1 = 0$, a contradiction. Thus $W = g_2^{k_2}W'$ with $W' \mid g_3^{k_3} \cdots g_l^{k_l}$.

Let $i \in [3, l]$ such that $g_i \mid W'$. We only need to show that $\text{ord}(g_i) = 2$. Assume to the contrary that $\text{ord}(g_i) > 2$. We set $X = Ug_i g_2^{-1}$, and then $|X| = |U| = D(G) - 1$. Suppose that X has a zero-sum subsequence T . Then $g_i^{k_i + 1} \mid T$ and by **A4** we obtain that $\text{ord}(g_i) > 2k_i$. Therefore, $g_i^{-k_i}T \mid U$ and $g_i^{-k_i}T \neq g_i^{-k_i}U$ but $\sigma(g_i^{-k_i}T) = -k_i g_i = \sigma(g_i^{-k_i}U)$ which implies that $2k_i g_i = 0$ by **A1**, a contradiction. Thus X is zero-sum free, and $|X| = D(G) - 1$ which imply that $\Sigma(X) = G \setminus \{0\}$. Therefore, X has a subsequence T such that $\sigma(T) = g_1 - g_2$.

Suppose that $g_i^{k_i + 1} \nmid T$. Then $T \mid U$, and by definition of X we have $g_2^{k_2} \nmid T$ which implies that $g_2T \mid U$. Since $\sigma(g_2T) = g_1 = \sigma(g_1)$, we obtain that $2g_1 = 0$ by **A1**, a contradiction.

Suppose that $g_i^{k_i + 1} \mid T$. Then $g_i^{-1}T \mid U$. Since $\sigma(g_i^{-1}T) = g_1 - g_2 - g_i = \sigma(g_i^{-1}W)$ and $g_i^{-1}T \neq g_i^{-1}W$, we obtain that $\text{supp}(g_i^{-1}T) \cap \text{supp}(g_i^{-1}W) = \emptyset$ and $U = g_i^{-1}T \cdot g_i^{-1}W$ by **A1**. Set $T_1 = g_i^{-1}T$, then $g_1^{k_1} g_i^{k_i} \mid T_1$ and $g_2 \nmid T_1$. It follows that $\sigma(g_1^{-1}g_2 T_1) = -g_i = \sigma(g_i^{-1}U)$ which implies that $2g_i = 0$ by **A1**, a contradiction. \square

Repeating the argument of **A5**, we can find another subsequence W_1 of U with $\sigma(W_1) = g_2 - g_1$ such that $W_1 = g_1^{k_1}W'_1$ and for any $h \mid W'_1$, $\text{ord}(h) = 2$. Set $Y = WW_1 \left(\prod_{h \in \text{supp}(W') \cap \text{supp}(W'_1)} h^2 \right)^{-1}$, then Y is a zero-sum subsequence of U . Since $\text{ord}(g_3) > 2$, we have $g_3 \nmid W$ and $g_3 \nmid W_1$ which imply that $g_3 \nmid Y$. It follows that Y is a proper zero-sum subsequence of U , a contradiction.

CASE 2.2: $|\{i \in [1, l] \mid \text{ord}(g_i) > 2\}| \leq 2$.

Since $k_1g_1 + \dots + k_lg_l = \sigma(U) = 0$ and $\text{ord}(k_1g_1) > 2$, it follows that $|\{i \in [1, l] \mid \text{ord}(g_i) > 2\}| = 2$, say, $\text{ord}(g_1) > 2$ and $\text{ord}(g_2) > 2$. Then **A4** implies that $\text{ord}(g_1) > 2k_1$ and $\text{ord}(g_2) > 2k_2$.

Suppose that $l = 2$. Then

$$D(G) - 1 = k_1 + k_2 \leq \frac{\text{ord}(g_1) - 1}{2} + \frac{\text{ord}(g_2) - 1}{2} \leq \exp(G) - 1$$

which implies that G is a cyclic group and $k_1 = k_2$. Since any minimal zero-sum sequence of length $|G| - 1$ over a cyclic group has the form $g^{|G|-2}(2g)$ for some generating element $g \in G$, it follows that $1 = k_2 = k_1$, and hence $|G| = 3$, a contradiction to $D(G) \geq 5$.

Suppose $l \geq 3$. Then $\exp(G)$ is even. We may assume that $\text{ord}(g_1) \geq \text{ord}(g_2)$. Since (g_3, \dots, g_l) is independent, we have $r(G) \geq l - 2$. Therefore,

$$\begin{aligned} \exp(G) + l - 4 &\leq D(G) - 1 = |U| = k_1 + k_2 + l - 2 \\ &\leq \lfloor \frac{\text{ord}(g_1) - 1}{2} \rfloor + \lfloor \frac{\text{ord}(g_2) - 1}{2} \rfloor + l - 2 \leq 2 \lfloor \frac{\text{ord}(g_1) - 1}{2} \rfloor + l - 2 \\ &\leq \text{ord}(g_1) + l - 3 \leq \exp(G) + l - 3. \end{aligned}$$

Since $\text{ord}(g_1) \mid \exp(G)$, we have that $\text{ord}(g_1) = \exp(G)$ is even. Thus $2 \lfloor \frac{\text{ord}(g_1) - 1}{2} \rfloor = \text{ord}(g_1) - 2$ which implies that $k_1 + k_2 = \lfloor \frac{\text{ord}(g_1) - 1}{2} \rfloor + \lfloor \frac{\text{ord}(g_2) - 1}{2} \rfloor = 2 \lfloor \frac{\text{ord}(g_1) - 1}{2} \rfloor$. Then $\text{ord}(g_1) = \text{ord}(g_2) = 2k_1 + 2 = 2k_2 + 2$. Since $\sigma(U) = k_1g_1 + k_2g_2 + g_3 + \dots + g_l = 0$, we have $2k_1g_1 + 2k_2g_2 = 0$ which implies that $2g_1 + 2g_2 = 0$. If $k_1 = k_2 \geq 2$, then $g_1^2g_2^2$ is an atom and divides UV , a contradiction. Thus $k_1 = k_2 = 1$, and hence $\text{ord}(g_1) = \text{ord}(g_2) = 4$. By **A3**, there exists a subsequence W of U such that $\sigma(W) = 2g_1$. If $g_1 \mid W$, then $g_1 \nmid g_1^{-1}W$ and $\sigma(g_1^{-1}W) = g_1 = \sigma(g_1)$ which implies that $2g_1 = 0$ by **A1**, a contradiction. If $g_1 \nmid W$, then $g_1 \nmid U(g_1W)^{-1}$ and $\sigma(U(g_1W)^{-1}) = g_1 = \sigma(g_1)$ which implies that $2g_1 = 0$ by **A1**, a contradiction. \square

4. PROOF OF THE MAIN RESULTS

In this final section we provide the proofs of all results presented in the Introduction (Theorem 1.1, Corollary 1.2, and Corollary 1.3).

Proof of Theorem 1.1 and of Corollary 1.2. Let H be a Krull monoid with finite class group G such that $|G| \geq 3$ and each class contains a prime divisor. Recall that the monoid of zero-sum sequences $\mathcal{B}(G)$ is a Krull monoid with class group isomorphic to G and each class contains a prime divisor. By Proposition 2.1, $\mathfrak{T}(H) = \mathfrak{T}(G)$ and $\mathfrak{c}(H) = \mathfrak{c}(G)$. Thus it is sufficient to prove Theorem 1.1 for the Krull monoid $\mathcal{B}(G)$.

Let \mathcal{O} be a holomorphy ring in a global field K , and R a classical maximal \mathcal{O} -order in a central simple algebra A over K such that every stably free left R -ideal is free. Then the monoid R^\bullet is a non-commutative Krull monoid ([15]), and all invariants under consideration of R^\bullet coincide with the respective invariants of a commutative Krull monoid whose class group is isomorphic to a ray class group of \mathcal{O} . These (highly non-trivial) transfer results are established in [31, 4], and are summarized in [4, Theorems 7.6 and 7.12]. Therefore, both for Theorem 1.1 and for Corollary 1.2, it is sufficient to prove the equivalence of the statements for a monoid of zero-sum sequences.

Let G be a finite abelian group with $|G| \geq 3$, and recall the inequalities

$$\mathfrak{T}(G) \leq \mathfrak{c}(G) \leq D(G).$$

(c) \Rightarrow (a) Suppose that G is isomorphic either to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$ or to $C_2 \oplus C_{2n}$ for some $n \geq 2$. Then Theorem **A** (in the Introduction) shows that $\mathfrak{c}(G) \leq D(G) - 1$. Since $D(C_2^{r-1} \oplus C_4) = r + 3$ and $D(C_2 \oplus C_{2n}) = 2n + 1$, Lemma 3.1.2 implies that $\mathfrak{c}(G) \geq \mathfrak{T}(G) \geq D(G) - 1$.

(a) \Rightarrow (b) Suppose that $\mathfrak{c}(G) = D(G) - 1$. By Theorem **A**, G is neither cyclic nor an elementary 2-group which implies that $D(G) \geq 5$. By Lemma 3.1.1, we have

$$D(G) - 1 = \mathfrak{c}(G) \leq \max \left\{ \left\lfloor \frac{1}{2} D(G) + 1 \right\rfloor, \mathfrak{T}(G) \right\} \leq D(G).$$

Thus, if $\mathfrak{T}(G) < D(G) - 1$, then $\left\lfloor \frac{1}{2}D(G) + 1 \right\rfloor \geq D(G) - 1$ which implies that $D(G) \leq 4$, a contradiction.

(b) \Rightarrow (c) Suppose that $\mathfrak{T}(G) = D(G) - 1$. Again by Theorem **A**, G is neither cyclic nor an elementary 2-group which implies that $D(G) \geq 5$. Then there exist $U, V \in \mathcal{A}(G)$ such that $\min(L(UV) \setminus \{2\}) = D(G) - 1$. Obviously, there are the following four cases (up to symmetry):

- $L(UV) = \{2, D(G) - 1, D(G)\}$.
- $L(UV) = \{2, D(G) - 1\}$ and $|U| = |V| = D(G)$.
- $L(UV) = \{2, D(G) - 1\}$ and $|U| - 1 = |V| = D(G)$.
- $L(UV) = \{2, D(G) - 1\}$ and $|U| = |V| = D(G) - 1$.

These cases are handled in the Propositions 3.5 to 3.8, and they imply that G is isomorphic either to $C_2^{r-1} \oplus C_4$ for some $r \geq 2$ or to $C_2 \oplus C_{2n}$ for some $n \geq 2$. \square

Proof of Corollary 1.3. Let G and G' be abelian groups such that $\mathcal{L}(G) = \mathcal{L}(G')$. Then

$$\mathfrak{T}(G) = \sup\{\min(L \setminus \{2\}) \mid 2 \in L \in \mathcal{L}(G)\} = \mathfrak{T}(G').$$

If G is finite, then $\mathfrak{T}(G) \leq c(G) \leq D(G) < \infty$. If G is infinite, then, by the Theorem of Kainrath (see [22] or [18, Section 7.3]), every finite set $L \subset \mathbb{N}_{\geq 2}$ lies in $\mathcal{L}(G)$, which implies that $\mathfrak{T}(G) = \infty$.

For $k \in \mathbb{N}$, we define the refined elasticities

$$\rho_k(G) = \sup\{\sup L \mid k \in L \in \mathcal{L}(G)\},$$

and observe that $\rho_k(G) = \rho_k(G')$. This implies that $kD(G) = \rho_{2k}(G) = \rho_{2k}(G') = kD(G')$ (see [18, Section 6.3]) for each $k \in \mathbb{N}$, and hence $D(G) = D(G')$.

Now suppose that $G' \in \{C_2^{r-1} \oplus C_4, C_2 \oplus C_{2n}\}$ where $r, n \geq 2$. Then Theorem 1.1 implies that $\mathfrak{T}(G') = D(G') - 1$. Since $\mathcal{L}(G) = \mathcal{L}(G')$, it follows that G is finite and that

$$\mathfrak{T}(G) = \mathfrak{T}(G') = D(G') - 1 = D(G) - 1,$$

whence Theorem 1.1 implies that $G \in \{C_2^{r-1} \oplus C_4, C_2 \oplus C_{2n}\}$ with $n, r \geq 2$. Suppose now that $n, r \geq 3$. Clearly, Condition (b) in Proposition 3.5 is equivalent to

$$(b') \quad \{2, D(G) - 1, D(G)\} \in \mathcal{L}(G).$$

Thus Proposition 3.5 implies in particular that $\mathcal{L}(C_2 \oplus C_{2n}) \neq \mathcal{L}(C_2^{r-1} \oplus C_4)$, and thus the assertion of Corollary 1.3 follows. \square

REFERENCES

- [1] D. Bachman, N. Baeth, and J. Gossell, *Factorizations of upper triangular matrices*, Linear Algebra Appl. **450** (2014), 138 – 157.
- [2] N.R. Baeth and A. Geroldinger, *Monoids of modules and arithmetic of direct-sum decompositions*, Pacific J. Math., to appear.
- [3] N.R. Baeth, V. Ponomarenko, D. Adams, R. Ardila, D. Hannasch, A. Kosh, H. McCarthy, and R. Rosenbaum, *Number theory of matrix semigroups*, Linear Algebra Appl. **434** (2011), 694 – 711.
- [4] N.R. Baeth and D. Smertnig, *Factorization theory in noncommutative settings*, <http://arxiv.org/abs/1402.4397>.
- [5] N.R. Baeth and R. Wiegand, *Factorization theory and decomposition of modules*, Am. Math. Mon. **120** (2013), 3 – 34.
- [6] P. Baginski, A. Geroldinger, D.J. Gryniewicz, and A. Philipp, *Products of two atoms in Krull monoids and arithmetical characterizations of class groups*, Eur. J. Comb. **34** (2013), 1244 – 1268.
- [7] Gyu Whan Chang and D. Smertnig, *Factorization in the self-idealization of a PID*, Boll. Unione Mat. Ital. **IX,6(2)** (2013), 363 – 377.
- [8] S.T. Chapman, P.A. García-Sánchez, and D. Llena, *The catenary and tame degree of numerical monoids*, Forum Math. **21** (2009), 117 – 129.
- [9] S.T. Chapman, P.A. García-Sánchez, D. Llena, V. Ponomarenko, and J.C. Rosales, *The catenary and tame degree in finitely generated commutative cancellative monoids*, Manuscr. Math. **120** (2006), 253 – 264.
- [10] S.T. Chapman, F. Gotti, and R. Pelayo, *On delta sets and their realizable subsets in Krull monoids with cyclic class groups*, Colloq. Math., to appear, –.
- [11] A. Facchini, *Krull monoids and their application in module theory*, Algebras, Rings and their Representations (A. Facchini, K. Fuller, C. M. Ringel, and C. Santa-Clara, eds.), World Scientific, 2006, pp. 53 – 71.

- [12] W. Gao, A. Geroldinger, and D.J. Gryniewicz, *Inverse zero-sum problems III*, Acta Arith. **141** (2010), 103 – 152.
- [13] P.A. García-Sánchez, I. Ojeda, and A. Sánchez-R.-Navarro, *Factorization invariants in half-factorial affine semigroups*, Internat. J. Algebra Comput. **23** (2013), 111 – 122.
- [14] A. Geroldinger, *Additive group theory and non-unique factorizations*, Combinatorial Number Theory and Additive Group Theory (A. Geroldinger and I. Ruzsa, eds.), Advanced Courses in Mathematics CRM Barcelona, Birkhäuser, 2009, pp. 1 – 86.
- [15] ———, *Non-commutative Krull monoids: a divisor theoretic approach and their arithmetic*, Osaka J. Math. **50** (2013), 503 – 539.
- [16] A. Geroldinger and D.J. Gryniewicz, *On the structure of minimal zero-sum sequences with maximal cross number*, J. Combinatorics and Number Theory **1** (2) (2009), 9 – 26.
- [17] A. Geroldinger, D.J. Gryniewicz, and W.A. Schmid, *The catenary degree of Krull monoids I*, J. Théor. Nombres Bordx. **23** (2011), 137 – 169.
- [18] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
- [19] A. Geroldinger and P. Yuan, *The set of distances in Krull monoids*, Bull. Lond. Math. Soc. **44** (2012), 1203 – 1208.
- [20] D.J. Gryniewicz, *Structural Additive Theory*, Developments in Mathematics, Springer, 2013.
- [21] F. Halter-Koch, *Ideal Systems. An Introduction to Multiplicative Ideal Theory*, Marcel Dekker, 1998.
- [22] F. Kainrath, *Factorization in Krull monoids with infinite class group*, Colloq. Math. **80** (1999), 23 – 30.
- [23] M. Omidali, *The catenary and tame degree of numerical monoids generated by generalized arithmetic sequences*, Forum Math. **24** (2012), 627 – 640.
- [24] A. Philipp, *A characterization of arithmetical invariants by the monoid of relations II: The monotone catenary degree and applications to semigroup rings*, Semigroup Forum, in print.
- [25] C. Reiher, *A proof of the theorem according to which every prime number possesses property B*, J. Lond. Math. Soc., to appear.
- [26] S. Savchev and F. Chen, *Long minimal zero-sum sequences in the groups $C_2^{r-1} \oplus C_{2k}$* , Integers **14** (2014), Paper A23.
- [27] W.A. Schmid, *Arithmetical characterization of class groups of the form $\mathbb{Z}/n\mathbb{Z} \oplus \mathbb{Z}/n\mathbb{Z}$ via the system of sets of lengths*, Abh. Math. Semin. Univ. Hamb. **79** (2009), 25 – 35.
- [28] ———, *Characterization of class groups of Krull monoids via their systems of sets of lengths: a status report*, Number Theory and Applications: Proceedings of the International Conferences on Number Theory and Cryptography (S.D. Adhikari and B. Ramakrishnan, eds.), Hindustan Book Agency, 2009, pp. 189 – 212.
- [29] ———, *Inverse zero-sum problems II*, Acta Arith. **143** (2010), 333 – 343.
- [30] ———, *The inverse problem associated to the Davenport constant for $C_2 \oplus C_2 \oplus C_{2n}$, and applications to the arithmetical characterization of class groups*, Electron. J. Comb. **18**(1) (2011), Research Paper 33.
- [31] D. Smertnig, *Sets of lengths in maximal orders in central simple algebras*, J. Algebra **390** (2013), 1 – 43.

UNIVERSITY OF GRAZ, INSTITUTE FOR MATHEMATICS AND SCIENTIFIC COMPUTING, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

E-mail address: alfred.geroldinger@uni-graz.at, qinghai.zhong@uni-graz.at